INDEX TO AUTHORS

AARONSON, STUART A. See STEPHENSON and AARONSON, 175

ADAMS, P. See VON BOEHMER, SHORTMAN, and ADAMS, 1648

ADLER, ANDY. See TAUB, ROSETT, ADLER, and MORSE, 1581

ALEXANDER, P. See EVANS, GRANT, COX, STEELE, and ALEXANDER, 1318

ALKAN, SEFIK S., BUSH, MAURICE E., NITECKI, DANUTE E., and GOODMAN, JOEL W. Antigen recognition and the immune response. Structural requirements in the side chain of tyrosine for immunogenicity of L-tyrosine-azobenzenearsonate, 387

—. See BUSH, ALKAN, NITECKI, and GOODMAN, 1478

ALTER, BARBARA J. See CLICK, BENCK, ALTER, and LOVCHIK, 1241

AMOS, BERNARD. See BERKE, SULLIVAN, and AMOS, 1594

AMSBAUGH, DIANA F., HANSEN, CARL T., PRESCOTT, BENJAMIN, STASHAK, PHILIP W., BARTHOLD, DAVID R., and BAKER, PHILLIP J. Genetic control of the antibody response to Type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness, 931

ANDERSON, TRAUTE-HEIDI, ROETHLE, JOAN, and AUERBACH, ROBERT. Induction of tolerance to sheep red blood cells, 1666

AOKI, TADAO. See HERBERMAN and AOKI, 94

AREND, WILLIAM P., and MANNIK, MART. In vitro adherence of soluble immune complexes to macrophages, 514

ASOFSKY, R. See LIEBERMAN, STIFFEL, ASOFSKY, MOUTON, BIOZZI, and BENACERRAF, 790

ATKINS, ELISHA. See CHUSID and ATKINS, 227

ATKINS, R. C. See ROWLEY, GOWANS, ATKINS, FORD, and SMITH, 499

AUERBACH, ROBERT. See ANDERSON, ROETHLE, and AUERBACH, 1666

AUSTEN, K. FRANK. See GOETZL and AUSTEN, 1564

—. See KALINER, ORANGE, and AUSTEN, 556

—. See KAPLAN and AUSTEN, 1378

BACH, FRITZ H., WIDMER, MICHAEL B., BACH, MARILYN L., and KLEIN, JAN. Serologically defined and lymphocyte-defined components of the major histocompatibility complex in the mouse, 1430

—. See WIDMER, SONDEL, and BACH, 398

BACH, MARILYN L. See BACH, WIDMER, BACH, and KLEIN, 1430

BAKER, PHILLIP J. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, BARTHOLD, and BAKER, 931

BARON, S. See WORTHINGTON, RABSON, and BARON, 277

BARTHOLD, DAVID R. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, BARTHOLD, and BAKER, 931

BASTEN, ANTONY. See FELDMANN and BASTEN, 49, 722

BATTISTO, J. R. See BUCSI, BOREK, and BATTISTO, 761

BEARN, ALEXANDER G. See DANES and BEARN, 1313

BEARN, ALEXANDER G. See DANES, HÜTTEROTH, CLEVE, and BEARN, 644
BENACERRAF, BARUJ. See DAVIE, PAUL, KATZ, and BENACERRAF, 426
—. See DUNHAM, UNANUE, and BENACERRAF, 403
—. See KATZ, HAMAOKA, and BENACERRAF, 1404
—. See LIEBERMAN, STIFFEL, ASOFSKY, MOUTON, BIOZZI, and BENACERRAF, 790
BENCK, LORETTA. See CLICK, BENCK, ALTER, and LOVCHIK, 1241
BENJAMINI, E. See SCIBIENSKI, FONG, and BENJAMINI, 1308
BENVENISTE, JACQUES, HENSON, PETER M., and COCHRANE, CHARLES G. Leukocyte-dependent histamine release from rabbit platelets. The role of IgE, basophils, and a platelet-activating factor, 1356
BERKE, GIDEON, SULLIVAN, KAREN A., and AMOS, BERNARD. Rejection of ascites tumor allografts. II. A pathway for cell-mediated tumor destruction in vitro by peritoneal exudate lymphoid cells, 1594
BERLIN, RICHARD D. See UKENA and BERLIN, 1
BERNSTEIN, DAVID. See BOKISCH, BERNSTEIN, and KRAUSE, 799
BIANCO, CELSO. See VITETTA, BIANCO, NUSSENZWEIG, and UHR, 81
BILLINGHAM, R. E. See ZAKARIAN and BILLINGHAM, 1545
BINZ, HANS, and LINDENMANN, JEAN. Cellular receptors: binding of radioactively labeled anti-alloantiserum, 872
BIOZZI, G. See LIEBERMAN, STIFFEL, ASOFSKY, MOUTON, BIOZZI, and BENACERRAF, 790
BLOOM, BARRY R. See MAILLARD and BLOOM, 185
BLUMENSTOCK, D. See RAPAPORT, WATANABE, CANNON, MOLLEN, BLUMENSTOCK, and FERREBEE, 1080
BOKISCH, VIKTOR A., BERNSTEIN, DAVID, and KRAUSE, RICHARD M. Occurrence of 19S and 7S anti-IgGs during hyperimmunization of rabbits with streptococci, 799
BOREK, F. See BUCSI, BOREK, and BATTISTO, 761
BOREL, YVES. See GOLAN and BOREL, 305
BORENSTEIN, DAVID. See SHIN, KALISS, BORENSTEIN, and GATELY, 375
BOWERS, WILLIAM E. Lysosomes in rat thoracic duct lymphocytes, 1394
BOYLE, W. See WAGNER, FELDMANN, BOYLE, and SCHRADER, 331
BRADLOW, H. LEON. See KAPPAS, BRADLOW, GILLETTE, and GALLAGHER, 1043
BRAI, MELCHIORRE, and OSLER, ABRAHAM G. Cobra venom-induced hemolysis. Activity levels in sera of patients with neoplastic and other diseases, 950
BRAIN, JOSEPH D. See GODLESKI and BRAIN, 630
BROOKS, WILLIAM H., NETSKY, MARTIN G., NORMANSELL, DAVID E., and HORWITZ, DAVID A. Depressed cell-mediated immunity in patients with primary intracranial tumors. Characterization of a humoral immunosuppressive factor, 1631
BUCSI, R. A., BOREK, F., and BATTISTO, J. R. Splenic replenishment of synergistic ability to bone marrow and thymic cells of neonatally splenectomized CBA mice, 761
BUSH, MAURICE E., ALKAN, SEFIK S., NITECKI, DANUTE E., and GOODMAN, JOEL W. Antigen recognition and the immune response. "Self-help" with symmetrical bifunctional antigen molecules, 1478
—. See ALKAN, BUSH, NITECKI, and GOODMAN, 387

CANNON, F. D. See RAPAPORT, WATANABE, CANNON, MOLLEN,

BLUMENSTOCK, and FERREBEE, 1080
CARP, RICHARD I., LICURSI, PAMELA C., MERZ, PATRICIA A., and MERZ, GEORGE S. Decreased percentage of polymorphonuclear neutrophils in mouse peripheral blood after inoculation with material from multiple sclerosis patients, 618
CELADA, FRANCO. See MACARIO, CONWAY DE MACARIO, FRANCESCHI, and CELADA, 353
CEROTTINI, JEAN-CHARLES. See GREY, KUBO, and CEROTTINI, 1323
CHEERS, CHRISTINA, and MILLER, J. F. A. P. Cell-to-cell interaction in the immune response. IX. Regulation of hapten-specific antibody class by carrier priming, 1661
CHEN, CHANG, and HIRSCH, JAMES G. The effects of mercaptoethanol and of peritoneal macrophages on the antibody-forming capacity of nonadherent mouse spleen cells in vitro, 604
CHOI, YONG SUNG, and GOOD, ROBERT A. Development of chicken lymphoid system. II. Synthesis of primordial immunoglobulin M by the bursa cells of chick embryo, 8
CHUSID, MICHAEL J., and ATKINS, ELISHA. Studies on the mechanism of penicillin-induced fever, 227
CLAMAN, HENRY N. See WALTERS, MOORHEAD, and CLAMAN, 546
CLEVE, HARTWIG, HAMAGUCHI, HIDEO, and HÜTTEROTH, THOMAS H. Preparation and further characterization of the MN glycoprotein of human erythrocyte membranes, 1140
—. See DANES, HÜTTEROTH, CLEVE, and BEARN, 644
CLICK, ROBERT E., BENCK, LORETTA, ALTER, BARBARA J., and LOVCHIK, JUDITH C. Immune responses in vitro. VI. Genetic control of the in vivo-in vitro discrepancies in 19S antibody synthesis, 1241
CLINTON, BRUCE A., and WEIGLE, WILLIAM O. Cellular events during the induction of experimental thyroiditis in the rabbit, 1605
COCHRANE, CHARLES G. See BENVENISTE, HENSON, and COCHRANE, 1356
COLIGAN, JOHN E. See TERRY, HENKART, COLIGAN, and TODD, 200
COLLEY, DANIEL G. See HAWIGER, MARNEY, COLLEY, and DES PREZ, 68
CONWAY DE MACARIO, EVERLY. See MACARIO, CONWAY DE MACARIO, FRANCESCHI, and CELADA, 353
COONS, ALBERT H. See RUBIN and COONS, 1501
CORNELIUS, EUGENE A. Rapid viral induction of murine lymphomas in the graft-*versus*-host reaction, 1533
COTRAN, RAMZI S. See GIMBRONE, LEAPMAN, COTRAN, and FOLKMAN, 261
COX, HELEN. See EVANS, GRANT, COX, STEELE, and ALEXANDER, 1318
CUDKOWICZ, G. See ITO, SHEARER, TRIZIO, and CUDKOWICZ, 381

DAHL, LEWIS K., LEITL, GEORGE, and HEINE, MARTHA. Influence of dietary potassium and sodium/potassium molar ratios on the development of salt hypertension, 318
DANES, B. SHANNON, and BEARN, ALEXANDER G. Oyster ciliary inhibition by cystic fibrosis culture medium, 1313
—, HÜTTEROTH, THOMAS H., CLEVE, HARTWIG, and BEARN, ALEXANDER G. The Hurler syndrome: a study of cultured lymphoid cell lines, 644
DAVIE, JOSEPH M., PAUL, WILLIAM E., KATZ, DAVID H., and BENACERRAF, BARUJ. Hapten-specific tolerance. Preferential depression of the high affinity antibody response, 426

DE DUVE, C. See PETERS, MÜLLER, and DE DUVE, 1117
DENNERT, GUNTHER, and TUCKER, DEREK F. Selective priming of T cells by chemically altered cell antigens, 656
DES PREZ, ROGER M. See HAWIGER, MARNEY, COLLEY, and DES PREZ, 68
DICKLER, H. B., and KUNKEL, H. G. Interaction of aggregated γ-globulin with B lymphocytes, 191
DIXON, FRANK J. See SCHMIDTKE and DIXON, 392
DOUGLAS, TOMMY C. Occurrence of a θ-like antigen in rats, 1054
DUBOS, RENÉ. See LEE and DUBOS, 1031
DUNHAM, EDWARD K., UNANUE, EMIL R., and BENACERRAF, BARUJ. Antigen binding and capping by lymphocytes of genetic nonresponder mice, 403
DUTTON, RICHARD W. Inhibitory and stimulatory effects of concanavalin A on the response of mouse spleen cell suspensions to antigen. I. Characterization of the inhibitory cell activity, 1445
DVORAK, HAROLD F. See HAMMOND and DVORAK, 1518

ECKNER, ROBERT J., and STEEVES, RICHARD A. A classification of the murine leukemia viruses. Neutralization of pseudotypes of Friend spleen focus-forming virus by type-specific murine antisera, 832
EDDLESTON, A. L. W. F. See LEU, EDDLESTON, HADDEN, and GOOD, 589
EIDINGER, DAVID, and GARRETT, THOMAS J. Studies of the regulatory effects of the sex hormones on antibody formation and stem cell differentiation, 1098
ESSNER, EDWARD. See OLIVER and ESSNER, 291
EVANS, R., GRANT, C. K., COX, HELEN, STEELE, KATHLEEN, and ALEXANDER, P. Thymus-derived lymphocytes produce an immunologically specific macrophage-arming factor, 1318
EYLAR, EDWIN H. See SPITLER, VON MULLER, FUDENBERG, and EYLAR, 156

FEIZI, T. See WERNET, FEIZI, and KUNKEL, 650
FELDMANN, MARC. Cell interactions in the immune response in vitro. V. Specific collaboration via complexes of antigen and thymus-derived cell immunoglobulin, 737
—. Induction of immunity and tolerance in vitro by hapten protein conjugates. III. Hapten inhibition studies of antigen binding to B cells in immunity and tolerance, 532
— and BASTEN, ANTONY. Cell interactions in the immune response in vitro. III. Specific collaboration across a cell-impermeable membrane, 49
— and —. IV. Comparison of the effects of antigen-specific and allogeneic thymus-derived cell factors, 722
—. See WAGNER, FELDMANN, BOYLE, and SCHRADER, 331
FENYÖ, EVA MARIA. See STROUK, GRUNDNER, FENYÖ, LAMON, SKURZAK, and KLEIN, 344
FERREBEE, J. W. See RAPAPORT, WATANABE, CANNON, MOLLEN, BLUMENSTOCK, and FERREBEE, 1080
FOLKMAN, JUDAH. See GIMBRONE, LEAPMAN, COTRAN, and FOLKMAN, 261
FONG, S. See SCIBIENSKI, FONG, and BENJAMINI, 1308
FORD, W. L. See ROWLEY, GOWANS, ATKINS, FORD, and SMITH, 499
FRANCESCHI, CLAUDIO. See MACARIO, CONWAY DE MACARIO, FRANCESCHI, and CELADA, 353

FRÖLAND, S. S., and NATVIG, J. B. Class, subclass, and allelic exclusion of membrane-bound Ig of human B lymphocytes, 409
FUDENBERG, H. HUGH. See SPITLER, VON MULLER, FUDENBERG, and EYLAR, 156

GALLAGHER, T. F. See KAPPAS, BRADLOW, GILLETTE, and GALLAGHER, 1043
GARRETT, THOMAS J. See EIDINGER and GARRETT, 1098
GATELY, MAURICE K. See SHIN, KALISS, BORENSTEIN, and GATELY, 375
GERSHON, RICHARD K., and LIEBHABER, STEPHEN A. The response of T cells to histocompatibility-2 antigens. Dose-response kinetics, 112
—. See GERY, GERSHON, and WAKSMAN, 128
GERY, IGAL, GERSHON, RICHARD K., and WAKSMAN, BYRON H. Potentiation of the T-lymphocyte response to mitogens. I. The responding cell, 128
— and WAKSMAN, BYRON H. Potentiation of the T-lymphocyte response to mitogens. II. The cellular source of potentiating mediator(s), 143
GILLETTE, PETER N. See KAPPAS, BRADLOW, GILLETTE, and GALLAGHER, 1043
GIMBRONE, MICHAEL A., JR., LEAPMAN, STEPHEN B., COTRAN, RAMZI S., and FOLKMAN, JUDAH. Tumor dormancy in vivo by prevention of neovascularization, 261
GINSBURG, HAIM. See HOLLANDER and GINSBURG, 1344
GODLESKI, JOHN J., and BRAIN, JOSEPH D. The origin of alveolar macrophages in mouse radiation chimeras, 630
GOETZL, EDWARD J., and AUSTEN, K. FRANK. A neutrophil-immobilizing factor derived from human leukocytes. I. Generation and partial characterization, 1564
GOLAN, DOV THEO, and BOREL, YVES. Deletion of hapten-binding cells by a highly radioactive ^{125}I conjugate, 305
GOLDHABER, PAUL. See TASHJIAN, VOELKEL, LEVINE, and GOLDHABER, 1329
GOLUB, EDWARD S. Brain-associated stem cell antigen: an antigen shared by brain and hemopoietic stem cells, 369
GOOD, ROBERT A. See CHOI and GOOD, 8
—. See LEU, EDDLESTON, HADDEN, and GOOD, 589
—. See TOIVANEN, TOIVANEN, and GOOD, 816
GOODMAN, JOEL W. See ALKAN, BUSH, NITECKI, and GOODMAN, 387
—. See BUSH, ALKAN, NITECKI, and GOODMAN, 1478
GOWANS, J. L. See ROWLEY, GOWANS, ATKINS, FORD, and SMITH, 499
GRANT, C. K. See EVANS, GRANT, COX, STEELE, and ALEXANDER, 1318
GREEN, IRA. See SHEVACH, PAUL, and GREEN, 1207
GREY, HOWARD M., KUBO, RALPH T., and CEROTTINI, JEAN-CHARLES. Thymus-derived (T) cell immunoglobulins. Presence of a receptor site for IgG and absence of large amounts of "buried" Ig determinants on T cells, 1323
GRIFFIN, DIANE E. See MCFARLAND, GRIFFIN, and JOHNSON, 216
GRUMET, F. CARL. See ORDAL and GRUMET, 1195
GRUNDNER, GERTRUD. See STROUK, GRUNDNER, FENYÖ, LAMON, SKURZAK, and KLEIN, 344

HADDEN, JOHN W. See LEU, EDDLESTON, HADDEN, and GOOD, 589
HAMAGUCHI, HIDEO. See CLEVE, HAMAGUCHI, and HÜTTEROTH, 1140
HAMAOKA, TOSHIYUKI. See KATZ,

HAMAOKA, and BENACERRAF, 1404
HAMMOND, M. ELIZABETH, and DVORAK, HAROLD F. Antigen-induced stimulation of glucosamine incorporation by guinea pig peritoneal macrophages in delayed hypersensitivity, 1518
HANSEN, CARL T. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, BARTHOLD, and BAKER, 931
HARTLEY, JANET W. See ROWE and HARTLEY, 1286
HAWIGER, JACEK, MARNEY, SAMUEL R., JR., COLLEY, DANIEL G., and DES PREZ, ROGER M. Complement-dependent platelet injury by staphylococcal protein A, 68
HEINE, MARTHA. See DAHL, LEITL, and HEINE, 318
HENKART, PIERRE A. See TERRY, HENKART, COLIGAN, and TODD, 200
HENSON, PETER M. See BENVENISTE, HENSON, and COCHRANE, 1356
HERBERMAN, RONALD B., and AOKI, TADAO. Immune and natural antibodies to syngeneic murine plasma cell tumors, 94
—. See ORTIZ DE LANDAZURI and HERBERMAN, 969
HERMAN, PETER G., YAMAMOTO, ITARU, and MELLINS, HARRY Z. Blood microcirculation in the lymph node during the primary immune response, 697
HIRSCH, JAMES G. See CHEN and HIRSCH, 604
—. See JONES and HIRSCH, 1173
—. See JONES, YEH, and HIRSCH, 1157
HOLLANDER, NURIT, and GINSBURG, HAIM. Specific adherence of in vitro differentiated lymphocytes to target cells, 1344
HOLM, G. See JONDAL, HOLM, and WIGZELL, 207
HORWITZ, DAVID A. See BROOKS, NETSKY, NORMANSELL, and HORWITZ, 1631
HÜTTEROTH, THOMAS H. See CLEVE, HAMAGUCHI, and HÜTTEROTH, 1140
—. See DANES, HÜTTEROTH, CLEVE, and BEARN, 644
HUMPHREY, WILLIAM, JR. See LIEBERMAN and HUMPHREY, 1222
—. See LIEBERMAN, PAUL, HUMPHREY, and STIMPFLING, 1231

ITO, T., SHEARER, G. M., TRIZIO, D., and CUDKOWICZ, G. B cell activation in vivo by non-antigen-specific interaction with T cells. Frequency of immune responses increased by immunization with two antigens, 381

JOHNSON, RICHARD T. See MCFARLAND, GRIFFIN, and JOHNSON, 216
JONDAL, M., HOLM, G., and WIGZELL, HANS. Surface markers on human T and B lymphocytes. I. A large population of lymphocytes forming nonimmune rosettes with sheep red blood cells, 207
JONES, THOMAS C., and HIRSCH, JAMES G. The interaction between *Toxoplasma gondii* and mammalian cells. II. The absence of lysosomal fusion with phagocytic vacuoles containing living parasites, 1173
—, YEH, SHIRLEY, and HIRSCH, JAMES G. The interaction between *Toxoplasma gondii* and mammalian cells. I. Mechanism of entry and intracellular fate of the parasite, 1157

KALINER, MICHAEL, ORANGE, ROBERT P., and AUSTEN, K. FRANK. Immunological release of histamine and slow-reacting substance of anaphylaxis from human lung. IV. Enhancement by cholinergic and α-adrenergic stimulation, 556
KALISS, NATHAN. See SHIN, KALISS, BORENSTEIN, and GATELY, 375

KAPLAN, ALLEN P., and AUSTEN, K. FRANK. The fibrinolytic pathway of human plasma. Isolation and characterization of the plasminogen proactivator, 1378

KAPPAS, ATTALLAH, BRADLOW, H. LEON, GILLETTE, PETER N., and GALLAGHER, T. F. Steroid studies in porphyria. I. A defect in the reductive transformation of natural steroid hormones in the hereditary liver disease, acute intermittent porphyria, 1043

KARNOVSKY, MORRIS J., UNANUE, EMIL R., and LEVENTHAL, MONIKA. Ligand-induced movement of lymphocyte membrane macromolecules. II. Mapping of surface moieties, 907

—. See RAVIOLA and KARNOVSKY, 466

—. See UNANUE, PERKINS, and KARNOVSKY, 885

KATZ, DAVID H., HAMAOKA, TOSHIYUKI, and BENACERRAF, BARUJ. Immunological tolerance in bone marrow-derived lymphocytes. I. Evidence for an intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404

— and OSBORNE, DAVID P., JR. The allogeneic effect in inbred mice. II. Establishment of the cellular interactions required for enhancement of antibody production by the graft-*versus*-host reaction, 455

—. See DAVIE, PAUL, KATZ, and BENACERRAF, 426

—. See OSBORNE and KATZ, 439

KLEIN, E. See LAMON, SKURZAK, KLEIN, and WIGZELL, 1072

KLEIN, GEORGE. See STROUK, GRUNDNER, FENYÖ, LAMON, SKURZAK, and KLEIN, 344

KLEIN, JAN. See BACH, WIDMER, BACH, and KLEIN, 1430

KLINMAN, NORMAN R. The mechanism of antigenic stimulation of primary and secondary clonal precursor cells, 241

KONDA, SUSUMU, NAKAO, YOSHINOBU, and SMITH, RICHARD T. Immunologic properties of mouse thymus cells. Identification of T cell functions within a minor, low-density subpopulation, 1461

KRAUSE, RICHARD M. See BOKISCH, BERNSTEIN, and KRAUSE, 799

KUBO, RALPH T. See GREY, KUBO, and CEROTTINI, 1323

KUNKEL, H. G. See DICKLER and KUNKEL, 191

—. See WERNET, FEIZI, and KUNKEL, 650

LAMELIN, J.-P., LISOWSKA-BERNSTEIN, B., MATTER, A., RYSER, J. E., and VASSALLI, P. Mouse thymus-independent and thymus-derived lymphoid cells. I. Immunofluorescent and functional studies, 984

—. See MATTER, LISOWSKA-BERNSTEIN, RYSER, LAMELIN, and VASSALLI, 1008

LAMON, ED W., SKURZAK, HENRYK M., KLEIN, E., and WIGZELL, HANS. In vitro cytotoxicity by a non-thymus-processed lymphocyte population with specificity for a virally determined tumor cell surface antigen, 1072

—. See STROUK, GRUNDNER, FENYÖ, LAMON, SKURZAK, and KLEIN, 344

LEAPMAN, STEPHEN B. See GIMBRONE, LEAPMAN, COTRAN, and FOLKMAN, 261

LEE, CHI-JEN, and DUBOS, RENÉ. Lasting biological effects of early environmental influences. VIII. Effects of neonatal infection, perinatal malnutrition, and crowding on catecholamine metabolism of brain, 1031

LEITL, GEORGE. See DAHL, LEITL, and HEINE, 318

LEU, RICHARD W., EDDLESTON, A. L. W. F., HADDEN, JOHN W., and GOOD, ROBERT A. Mechanism of action of migration inhibitory factor

(MIF). I. Evidence for a receptor for MIF present on the peritoneal macrophage but not on the alveolar macrophage, 589
LEVENTHAL, MONIKA. See KARNOVSKY, UNANUE, and LEVENTHAL, 907
LEVINE, LAWRENCE. See TASHJIAN, VOELKEL, LEVINE, and GOLDHABER, 1329
LICURSI, PAMELA C. See CARP, LICURSI, MERZ, and MERZ, 618
LIEBERMAN, ROSE, and HUMPHREY, WILLIAM, JR. Association of H-2 types with genetic control of immune responsiveness to IgG (γ2a) allotypes in the mouse, 1222
—, PAUL, WILLIAM E., HUMPHREY, WILLIAM, JR., and STIMPFLING, J. H. H-2-linked immune response (Ir) genes. Independent loci for *Ir-IgG* and *Ir-IgA* genes, 1231
—, STIFFEL, C., ASOFSKY, R., MOUTON, D., BIOZZI, G., and BENACERRAF, BARUJ. Genetic factors controlling anti-sheep erythrocyte antibody response and immunoglobulin synthesis in backcross and F_2 progeny of mice genetically selected for "high" or "low" antibody synthesis, 790
LIEBHABER, STEPHEN A. See GERSHON and LIEBHABER, 112
LINDENMANN, JEAN. See BINZ and LINDENMANN, 872
LISOWSKA-BERNSTEIN, B. See LAMELIN, LISOWSKA-BERNSTEIN, MATTER, RYSER, and VASSALLI, 984
—. See MATTER, LISOWSKA-BERNSTEIN, RYSER, LAMELIN, and VASSALLI, 1008
LOVCHIK, JUDITH C. See CLICK, BENCK, ALTER, and LOVCHIK, 1241

MACARIO, ALBERTO J. L., CONWAY DE MACARIO, EVERLY, FRANCESCHI, CLAUDIO, and CELADA, FRANCO. Maturation of the immune response in vitro. Focal fluctuation and changes in affinity of anti-β-D-galactosidase-activating antibody, 353
MAILLARD, JEAN, and BLOOM, BARRY R. Immunological adjuvants and the mechanism of cell cooperation, 185
MANNIK, MART. See AREND and MANNIK, 514
MARNEY, SAMUEL R., JR. See HAWIGER, MARNEY, COLLEY, and DES PREZ, 68
MATTER, A., LISOWSKA-BERNSTEIN, B., RYSER, J. E., LAMELIN, J.-P., and VASSALLI, P. Mouse thymus-independent and thymus-derived lymphoid cells. II. Ultrastructural studies, 1008
—. See LAMELIN, LISOWSKA-BERNSTEIN, MATTER, RYSER, and VASSALLI, 984
MCFARLAND, HENRY F., GRIFFIN, DIANE E., and JOHNSON, RICHARD T. Specificity of the inflammatory response in viral encephalitis. I. Adoptive immunization of immunosuppressed mice infected with Sindbis virus, 216
MCKENZIE, IAN F. C. See SILVER, MCKENZIE, and WINN, 1063
MELLINS, HARRY Z. See HERMAN, YAMAMOTO, and MELLINS, 697
MELMON, KENNETH L. See SHEARER, MELMON, WEINSTEIN, and SELA, 1302
MERZ, GEORGE S. See CARP, LICURSI, MERZ, and MERZ, 618
MERZ, PATRICIA A. See CARP, LICURSI, MERZ, and MERZ, 618
MILES, R. E. Probability analysis. Appendix to Schoefl, 568
MILLER, J. F. A. P. See CHEERS and MILLER, 1661
MOLLEN, N. See RAPAPORT, WATANABE, CANNON, MOLLEN, BLUMENSTOCK, and FERREBEE, 1080
MOND, J. J., TAKAHASHI, T., and THORBECKE, G. J. Surface antigens of immunocompetent cells. III. In vitro studies of the role of

B and T cells in immunological memory, 663
MOND, J. J., TAKAHASHI, T., and THORBECKE, G. J. Thymus-derived cell (T cell) activation by heterologous antigens as a replacement of specific immune T cells in the transfer of the secondary response to sheep erythrocytes, 715
MOORHEAD, JOHN W. See WALTERS, MOORHEAD, and CLAMAN, 546
MORSE, STEPHEN I. See TAUB, ROSETT, ADLER, and MORSE, 1581
MOSIER, DONALD E., and PIERCE, CARL W. Functional maturation of thymic lymphocyte populations in vitro, 1484
MOUTON, D. See LIEBERMAN, STIFFEL, ASOFSKY, MOUTON, BIOZZI, and BENACERRAF, 790
MÜLLER, M. See PETERS, MÜLLER, and DE DUVE, 1117

NAKAO, YOSHINOBU. See KONDA, NAKAO, and SMITH, 1461
NATVIG, J. B. See FRÖLAND and NATVIG, 409
NETSKY, MARTIN G. See BROOKS, NETSKY, NORMANSELL, and HORWITZ, 1631
NIELSEN, H. E. Reactivity of lymphocytes from germfree rats in mixed leukocyte culture and in graft-*versus*-host reaction, 417
NITECKI, DANUTE E. See ALKAN, BUSH, NITECKI, and GOODMAN, 387
—. See BUSH, ALKAN, NITECKI, and GOODMAN, 1478
NORMANSELL, DAVID E. See BROOKS, NETSKY, NORMANSELL, and HORWITZ, 1631
NUSSENZWEIG, VICTOR. See VITETTA, BIANCO, NUSSENZWEIG, and UHR, 81

O'BRIEN, JOHN S. See ZIELKE, VEATH, and O'BRIEN, 197
OLIVER, CONSTANCE, and ESSNER, EDWARD. Protein transport in mouse kidney utilizing tyrosinase as an ultrastructural tracer, 291
ORANGE, ROBERT P. See KALINER, ORANGE, and AUSTEN, 556
ORDAL, JOHN C., and GRUMET, F. CARL. Genetic control of the immune response. The effect of graft-*versus*-host reaction on the antibody response to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys in non-responder mice, 1195
ORTIZ DE LANDAZURI, MANUEL, and HERBERMAN, RONALD B. In vitro activation of cellular immune response to Gross virus-induced lymphoma, 969
OSBORNE, DAVID P., JR., and KATZ, DAVID H. The allogeneic effect in inbred mice. I. Experimental conditions for the enhancement of hapten-specific secondary antibody responses by the graft-*versus*-host reaction, 439
—. See KATZ and OSBORNE, 455
OSLER, ABRAHAM G. See BRAI and OSLER, 950

PAUL, WILLIAM E. See DAVIE, PAUL, KATZ, and BENACERRAF, 426
—. See LIEBERMAN, PAUL, HUMPHREY, and STIMPFLING, 1231
—. See SHEVACH, PAUL, and GREEN, 1207
PERKINS, WILLIAM D. See UNANUE, PERKINS, and KARNOVSKY, 885
PETERS, T. J., MÜLLER, M., and DE DUVE, C. Lysosomes of the arterial wall. I. Isolation and subcellular fractionation of cells from normal rabbit aorta, 1117
PIERCE, CARL W. See MOSIER and PIERCE, 1484
PIGUET, P.-F., and VASSALLI, P. Thymus-independent (B) cell proliferation in spleen cell cultures of mouse radiation chimeras stimulated by phytohemagglutinin or allogeneic cells, 962

PRESCOTT, BENJAMIN. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, BARTHOLD, and BAKER, 931

RABSON, A. S. See WORTHINGTON, RABSON, and BARON, 277

RAPAPORT, F. T., WATANABE, K., CANNON, F. D., MOLLEN, N., BLUMENSTOCK, D., and FERREBEE, J. W. Histocompatibility studies in a closely bred colony of dogs. IV. Tolerance to bone marrow, kidney, and skin allografts in DL-A-identical radiation chimeras, 1080

RAVIOLA, ELIO, and KARNOVSKY, MORRIS J. Evidence for a blood-thymus barrier using electron-opaque tracers, 466

ROETHLE, JOAN. See ANDERSON, ROETHLE, and AUERBACH, 1666

ROSETT, WALTER. See TAUB, ROSETT, ADLER, and MORSE, 1581

ROSS, RUSSELL. See STEMERMAN and ROSS, 769

ROWE, WALLACE P. Studies of genetic transmission of murine leukemia virus by AKR mice. I. Crosses with *Fv-1*^{n} strains of mice, 1272

— and HARTLEY, JANET W. Studies of genetic transmission of murine leukemia virus by AKR mice. II. Crosses with *Fv-1*^{b} strains of mice, 1286

ROWLEY, D. A., GOWANS, J. L., ATKINS, R. C., FORD, W. L., and SMITH, MARILYN E. The specific selection of recirculating lymphocytes by antigen in normal and preimmunized rats, 499

RUBIN, ARNOLD S., and COONS, ALBERT H. Specific heterologous enhancement of immune responses. IV. Specific generation of a thymus-derived enhancing factor, 1501

RYSER, J. E. See LAMELIN, LISOWSKA-BERNSTEIN, MATTER, RYSER, and VASSALLI, 984

RYSER, J. E. See MATTER, LISOWSKA-BERNSTEIN, RYSER, LAMELIN, and VASSALLI, 1008

SABBADINI, E. See SINGH, SABBADINI, and SEHON, 39

SCHIRRMACHER, VOLKER, and WIGZELL, HANS. Immune responses against native and chemically modified albumins in mice. I. Analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against methylated bovine serum albumin, 1616

SCHMIDTKE, JON R., and DIXON, FRANK J. Immune response to a hapten coupled to a nonimmunogenic carrier. Influence of lipopolysaccharide, 392

SCHOEFL, G. I. The migration of lymphocytes across the vascular endothelium in lymphoid tissue. A reexamination. With an Appendix by R. E. Miles. Probability analysis, 568

SCHRADER, J. W. See WAGNER, FELDMANN, BOYLE, and SCHRADER, 331

SCIBIENSKI, R., FONG, S., and BENJAMINI, E. Cross-tolerance between serologically non-cross-reacting forms of egg white lysozyme, 1308

SEHON, A. H. See SINGH, SABBADINI, and SEHON, 39

SELA, MICHAEL. See SHEARER, MELMON, WEINSTEIN, and SELA, 1302

SHEARER, G. M., MELMON, KENNETH L., WEINSTEIN, YACOB, and SELA, MICHAEL. Regulation of antibody response by cells expressing histamine receptors, 1302

—. See ITO, SHEARER, TRIZIO, and CUDKOWICZ, 381

SHEVACH, ETHAN M., PAUL, WILLIAM E., and GREEN, IRA. Histocompatibility-linked immune response gene function in guinea pigs. Specific inhibition of antigen-induced

lymphocyte proliferation by alloantisera, 1207

SHIN, HYUN S., KALISS, NATHAN, BORENSTEIN, DAVID, and GATELY, MAURICE K. Antibody-mediated suppression of grafted lymphoma cells. II. Participation of macrophages, 375

SHORTMAN, K. See VON BOEHMER, SHORTMAN, and ADAMS, 1648

SILVER, DONALD M., MCKENZIE, IAN F. C., and WINN, HENRY J. Variations in the responses of C57BL/10J and A/J mice to sheep red blood cells. I. Serological characterization and genetic analysis, 1063

SINGH, JAGAT N., SABBADINI, E., and SEHON, A. H. Cytotoxicity in graft-*versus*-host reaction. I. Role of donor and host spleen cells, 39

SKURZAK, HENRYK M. See LAMON, SKURZAK, KLEIN, and WIGZELL, 1072

—. See STROUK, GRUNDNER, FENYÖ, LAMON, SKURZAK, and KLEIN, 344

SMITH, MARILYN E. See ROWLEY, GOWANS, ATKINS, FORD, and SMITH, 499

SMITH, RICHARD T. See KONDA, NAKAO, and SMITH, 1461

SONDEL, PAUL M. See WIDMER, SONDEL, and BACH, 398

SPITLER, LYNN E., VON MULLER, CHRISTINE M., FUDENBERG, H. HUGH, and EYLAR, EDWIN H. Experimental allergic encephalitis. Dissociation of cellular immunity to brain protein and disease production, 156

STASHAK, PHILIP W. See AMSBAUGH, HANSEN, PRESCOTT, STASHAK, BARTHOLD, and BAKER, 931

STEELE, KATHLEEN. See EVANS, GRANT, COX, STEELE, and ALEXANDER, 1318

STEEVES, RICHARD A. See ECKNER and STEEVES, 832

STEMERMAN, MICHAEL B., and ROSS, RUSSELL. Experimental arteriosclerosis. I. A fibrous plaque formation in primates, an electron microscope study, 769

STEPHENSON, JOHN R., and AARONSON, STUART A. Genetic factors influencing C-type RNA virus induction, 175

STIFFEL, C. See LIEBERMAN, STIFFEL, ASOFSKY, MOUTON, BIOZZI, and BENACERRAF, 790

STIMPFLING, J. H. See LIEBERMAN, PAUL, HUMPHREY, and STIMPFLING, 1231

STROBER, SAMUEL. Initiation of antibody responses by different classes of lymphocytes. V. Fundamental changes in the physiological characteristics of virgin thymus-independent ("B") lymphocytes and "B" memory cells, 851

STROUK, VLADIMIR, GRUNDNER, GERTRUD, FENYÖ, EVA MARIA, LAMON, ED W., SKURZAK, HENRYK M., and KLEIN, GEORGE. Lack of distinctive surface antigen on cells transformed by murine sarcoma virus, 344

SULLIVAN, KAREN A. See BERKE, SULLIVAN, and AMOS, 1594

SWANSON, JOHN. Studies on gonococcus infection. II. Freeze-fracture, freeze-etch studies on gonococci, 1258

TAKAHASHI, T. See MOND, TAKAHASHI, and THORBECKE, 663, 715

TASHJIAN, ARMEN H., JR., VOELKEL, EDWARD F., LEVINE, LAWRENCE, and GOLDHABER, PAUL. Evidence that the bone resorption-stimulating factor produced by mouse fibrosarcoma cells is prostaglandin E_2. A new model for the hypercalcemia of cancer, 1329

TAUB, ROBERT N., ROSETT, WALTER, ADLER, ANDY, and MORSE, STEPHEN I. Distribution of labeled lymph node cells in mice during the lymphocytosis induced by *Bordetella pertussis*, 1581

TERRY, WILLIAM D., HENKART, PIERRE A., COLIGAN, JOHN E., and TODD, CHARLES W. Structural studies of the major glycoprotein in preparations with carcinoembryonic antigen activity, 200
THORBECKE, G. J. See MOND, TAKAHASHI, and THORBECKE, 663, 715
TODD, CHARLES W. See TERRY, HENKART, COLIGAN, and TODD, 200
TOIVANEN, AULI. See TOIVANEN, TOIVANEN, and GOOD, 816
TOIVANEN, PAAVO, TOIVANEN, AULI, and GOOD, ROBERT A. Ontogeny of bursal function in chicken. III. Immunocompetent cell for humoral immunity, 816
TRIZIO, D. See ITO, SHEARER, TRIZIO, and CUDKOWICZ, 381
TUCKER, DEREK F. See DENNERT and TUCKER, 656

UHR, JONATHAN W. See VITETTA, BIANCO, NUSSENZWEIG, and UHR, 81
—. See VITETTA and UHR, 676
UKENA, THOMAS E., and BERLIN, RICHARD D. Effect of colchicine and vinblastine on the topographical separation of membrane functions, 1
UNANUE, EMIL R., PERKINS, WILLIAM D., and KARNOVSKY, MORRIS J. Ligand-induced movement of lymphocyte membrane macromolecules. I. Analysis by immunofluorescence and ultrastructural radioautography, 885
—. See DUNHAM, UNANUE, and BENACERRAF, 403
—. See KARNOVSKY, UNANUE, and LEVENTHAL, 907

VASSALLI, P. See LAMELIN, LISOWSKA-BERNSTEIN, MATTER, RYSER, and VASSALLI, 984
—. See MATTER, LISOWSKA-BERNSTEIN, RYSER, LAMELIN, and VASSALLI, 1008
—. See PIGUET and VASSALLI, 962
VEATH, M. LOIS. See ZIELKE, VEATH, and O'BRIEN, 197
VITETTA, ELLEN S., BIANCO, CELSO, NUSSENZWEIG, VICTOR, and UHR, JONATHAN W. Cell surface immunoglobulin. IV. Distribution among thymocytes, bone marrow cells, and their derived populations, 81
— and UHR, JONATHAN W. Cell surface immunoglobulin. V. Release from murine splenic lymphocytes, 676
VOELKEL, EDWARD F. See TASHJIAN, VOELKEL, LEVINE, and GOLDHABER, 1329
VOLKMAN, ALVIN. The host cell response in the local graft-*versus*-host reaction induced in the kidneys of F_1 rats by parental thoracic duct lymphocytes, 21
VON BOEHMER, H., SHORTMAN, K., and ADAMS, P. Nature of the stimulating cell in the syngeneic and the allogeneic mixed lymphocyte reaction in mice, 1684
VON MULLER, CHRISTINE M. See SPITLER, VON MULLER, FUDENBERG, and EYLAR, 156

WAGNER, HERMANN, FELDMANN, MARC, BOYLE, W., and SCHRADER, J. W. Cell-mediated immune response in vitro. III. The requirement for macrophages in cytotoxic reactions against cell-bound and subcellular alloantigens, 331
WAKSMAN, BYRON H. See GERY, GERSHON, and WAKSMAN, 128
—. See GERY and WAKSMAN, 143
—. See YOSHINAGA, YOSHINAGA, and WAKSMAN, 956
WALTERS, CURLA S., MOORHEAD, JOHN W., and CLAMAN, HENRY N. Immunity and tolerance to a hapten (NIP) coupled to an isologous carrier (mouse γ-globulin), 546
WATANABE, K. See RAPAPORT, WATANABE, CANNON, MOLLEN,

BLUMENSTOCK, and FERREBEE, 1080
WEIGLE, WILLIAM O. See CLINTON and WEIGLE, 1605
WEINSTEIN, YACOB. See SHEARER, MELMON, WEINSTEIN, and SELA, 1302
WERNET, P., FEIZI, T., and KUNKEL, H. G. Idiotypic determinants of immunoglobulin M detected on the surface of human lymphocytes by cytotoxicity assays, 650
WIDMER, MICHAEL B., SONDEL, PAUL M., and BACH, FRITZ H. Responsiveness of germfree mice in mixed leukocyte culture. Reexamining the nature of the responding unit, 398
—. See BACH, WIDMER, BACH, and KLEIN, 1430
WIGZELL, HANS. See JONDAL, HOLM, and WIGZELL, 207
—. See LAMON, SKURZAK, KLEIN, and WIGZELL, 1072
—. See SCHIRRMACHER and WIGZELL, 1616
WINN, HENRY J. See SILVER, MCKENZIE, and WINN, 1063
WORTHINGTON, M., RABSON, A. S., and BARON, S. Mechanism of recovery from systemic vaccinia virus infection. I. The effects of cyclophosphamide, 277

YAMAMOTO, ITARU. See HERMAN, YAMAMOTO, and MELLINS, 697
YEH, SHIRLEY. See JONES, YEH, and HIRSCH, 1157
YOSHINAGA, AIKO. See YOSHINAGA, YOSHINAGA, and WAKSMAN, 956
YOSHINAGA, MASARU, YOSHINAGA, AIKO, and WAKSMAN, BYRON H. Regulation of lymphocyte responses in vitro. I. Regulatory effect of macrophages and thymus-dependent (T) cells on the response of thymus-independent (B) lymphocytes to endotoxin, 956

ZAKARIAN, SIRAIK, and BILLINGHAM, R. E. Studies on normal and immune lymphocyte transfer reactions in guinea pigs, with special reference to the cellular contribution of the host, 1545
ZIELKE, KLAUS, VEATH, M. LOIS, and O'BRIEN, JOHN S. Fucosidosis: Deficiency of alpha-L-fucosidase in cultured skin fibroblasts, 197

Statement of Ownership, Management, and Circulation required by Act of 23 October 1962; Section 4369, Title 39, United States Code: The Journal of Experimental Medicine. Published monthly at Mount Royal and Guilford Avenues, Baltimore, Maryland 21202, for The Rockefeller University Press, New York 10021. Editors: René Dubos, Henry G. Kunkel, Maclyn McCarty, The Rockefeller University, New York 10021. Owned by The Rockefeller University, New York 10021, nonprofit corporation, authorized to mail at special rates. Purpose, function, and nonprofit status of the organization and the exempt status for Federal income tax purposes have not changed during the preceding 12 months. Following statistics represent 12 month average and number nearest to date of filing: copies printed for each issue, 3639; sales through dealers, street vendors, counter sales, 0; mail subscriptions, 3073. Total paid circulation, 3163; free distribution, 90; office use, left-over, unaccounted, spoiled, 476; net press run, 3639. Single issue nearest filing date June 1972. (Signed W. A. Bayless, Director.)

INDEX TO SUBJECTS

ACID, ribonucleic, C-type, virus induction, genetic factors influencing, 175
Adjuvants, immunological, and cell cooperation mechanism, 185
α-Adrenergic and cholinergic stimulation, enhancement of immunological release of histamine and slow-reacting substance of anaphylaxis from human lung by, 556
Albumin, serum, bovine, methylated, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against, 1616
Albumins, native and chemically modified, immune responses against, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against methylated bovine serum albumin, 1616
Allelic exclusion, class, and subclass of membrane-bound Ig of human bone marrow-derived lymphocytes, 409
Allergic encephalitis, dissociation of cellular immunity to brain protein and disease production, 156
Alloantigens, cell-bound and subcellular, requirement for macrophages in cytotoxic reactions against, 331
Alloantisera, specific inhibition of antigen-induced lymphocyte proliferation by, histocompatibility-linked immune response gene function, 1207
Alloantiserum, anti-, radioactively labeled, binding of, 872
Allogeneic cells or phytohemagglutinin, thymus-independent (B) cell proliferation in spleen cell cultures of radiation chimeras stimulated by, 962
— effect in inbred mice, conditions for enhancement of hapten-specific secondary antibody responses by graft-*versus*-host reaction, 439
Allogeneic effect in inbred mice, establishment of cellular interactions required for antibody production enhancement by graft-*versus*-host reaction, 455
— and syngeneic mixed lymphocyte reaction, nature of stimulating cell in, 1648
— thymus-derived and antigenic-specific cell factors, comparison of effects in immune response in vitro, 722
Allografts, skin, bone marrow, and kidney in DL-A-identical radiation chimeras, tolerance to, histocompatibility studies in closely bred colony of dogs, 1080
—, tumor, ascites, rejection, pathway for cell-mediated tumor destruction in vitro by peritoneal exudate lymphoid cells, 1594
Allotypes, IgG (γ2a) in mouse, association of H-2 types with genetic control of immune responsiveness to, 1222
Alveolar macrophage, evidence for receptor for migration inhibitory factor (MIF) present on peritoneal macrophage but not on, 589
— macrophages, origin in radiation chimeras, 630
Anaphylaxis, slow-reacting substance of, and histamine, immunological release from human lung, enhancement by cholinergic and α-adrenergic stimulation, 556
Antibodies, immune and natural, to syngeneic murine plasma cell tumors, 94
Antibody, anti-β-D-galactosidase-ac-

tivating, focal fluctuation and changes in affinity of, 353
Antibody class, hapten-specific, regulation by carrier priming, cell-to-cell interaction in immune response, 1661
— formation and stem cell differentiation, regulatory effects of sex hormones on, 1098
— production, enhancement by graft-*versus*-host reaction, establishment of cellular interactions required for, 455
— response by cells expressing histamine receptors, regulation of, 1302
— response, erythrocyte, anti-sheep, and immunoglobulin synthesis in backcross and F_2 progeny of mice genetically selected for high or low antibody synthesis, genetic factors controlling, 790
— response, high affinity, preferential depression of, hapten-specific tolerance, 426
— response to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys, effect of graft-*versus*-host reaction, genetic control of immune response, 1195
— response to Type III pneumococcal polysaccharide, genetic control of, evidence that X-linked gene plays decisive role in determining responsiveness, 931
— responses, hapten-specific secondary, conditions for enhancement by graft-*versus*-host reaction, 439
— responses, initiation by different classes of lymphocytes, fundamental changes in physiological characteristics of virgin thymus-independent (B) lymphocytes and B memory cells, 851
— synthesis, 19S, genetic control of in vivo-in vitro discrepancies in, immune responses in vitro, 1241
Antibody-forming capacity of nonadherent spleen cells in vitro, effects of mercaptoethanol and peritoneal macrophages on, 604
Antibody-forming cells, evidence for intracellular mechanism of inactivation of hapten-specific precursors of, immunological tolerance in bone marrow-derived lymphocytes, 1404
Antibody-mediated suppression of grafted lymphoma cells, participation of macrophages, 375
Anti-β-D-galactosidase-activating antibody, focal fluctuation and changes in affinity of, 353
Antigen activity, carcinoembryonic, structural studies of major glycoprotein in preparations with, 200
— binding and capping by lymphocytes of genetic nonresponder mice, 403
— binding to non-thymus-dependent cells in immunity and tolerance, hapten inhibition studies of, 532
—, inhibitory and stimulatory effects of concanavalin A on response of spleen cell suspensions to, characterization of inhibitory cell activity, 1445
—, θ-like, occurrence in rats, 1054
— recognition and immune response, self-help with symmetrical bifunctional antigen molecules, 1478
— recognition and immune response, structural requirements in tyrosine side chain for immunogenicity of L-tyrosine-azobenzenearsonate, 387
—, specific selection of recirculating lymphocytes by, 499
—, stem cell, brain-associated, shared by brain and hemopoietic stem cells, 369
—, surface, distinctive, lack on cells transformed by murine sarcoma virus, 344
—, surface, tumor cell, virally determined, in vitro cytotoxicity by non-thymus-processed lymphocyte population with specificity for, 1072

Antigen and thymus-derived cell immunoglobulin complexes, specific collaboration via in immune response in vitro, 737
Antigenic stimulation of primary and secondary clonal precursor cells, mechanism, 241
Antigen-induced lymphocyte proliferation, specific inhibition by alloantisera, histocompatibility-linked immune response gene function, 1207
— stimulation of glucosamine incorporation by peritoneal macrophages in delayed hypersensitivity, 1518
Antigen-specific and allogeneic thymus-derived cell factors, comparison of effects in immune response in vitro, 722
—, non-, interaction with thymus-derived cells, bone marrow-derived cell activation in vivo by, frequency of immune responses increased by immunization with two antigens, 381
Antigens, cell, chemically altered, selective priming of thymus-derived cells by, 656
—, heterologous, thymus-derived, cell activation by as replacement of specific immune T cells in transfer of secondary response to erythrocytes, 715
—, histocompatibility-2, response of thymus-influenced cells to, dose-response kinetics, 112
—, surface, of immunocompetent cells, in vitro studies of role of bone marrow-derived and thymus-derived cells in immunological memory, 663
—, two, frequency of immune responses increased by immunization with, bone marrow-derived cell activation in vivo by non-antigen-specific interaction with thymus-derived cells, 381
Antisera, murine, type-specific, neutralization of pseudotypes of Friend spleen focus-forming virus by, 832
Aorta, isolation and subcellular fractionation of cells from, lysosomes of arterial wall, 1117
Arterial wall, lysosomes, isolation and subcellular fractionation of cells from aorta, 1117
Arteriosclerosis, fibrous plaque formation in primates, electron microscope study, 769
Ascites tumor allografts, rejection, pathway for cell-mediated tumor destruction in vitro by peritoneal exudate lymphoid cells, 1594
Azobenzenearsonate, L-tyrosine-, structural requirements in side chain of tyrosine for immunogenicity of, 387

BASOPHILS, IgE, and platelet-activating factor, role in leukocyte-dependent histamine release from platelets, 1356
Biological effects, lasting, of early environmental influences, effects of neonatal infection, perinatal malnutrition, and crowding on catecholamine metabolism of brain, 1031
Blood cells, red. *See* Erythrocytes.
— microcirculation in lymph node during primary immune response, 697
—, peripheral, decreased percentage of polymorphonuclear neutrophils in after inoculation with material from multiple sclerosis patients, 618
Blood-thymus barrier, evidence for using electron-opaque tracers, 466
Bone marrow cells, thymocytes, and derived populations, cell surface immunoglobulin distribution among, 81
— marrow, kidney, and skin allografts in DL-A-identical radiation chimeras, tolerance to, histocompatibility studies in closely bred colony of dogs, 1080
— marrow and thymic cells of neo-

natally splenectomized CBA mice, splenic replenishment of synergistic ability to, 761
Bone marrow-derived cell activation in vivo by non-antigen-specific interaction with thymus-derived cells, frequency of immune responses increased by immunization with two antigens, 381
— marrow-derived lymphocytes, human, class, subclass, and allelic exclusion of membrane-bound Ig of, 409
— marrow-derived lymphocytes, immunological tolerance in, evidence for intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404
— marrow-derived and thymus-derived cells, role in immunological memory, in vitro studies, 663
— marrow-derived and thymus-derived lymphocytes, human, surface markers on, large population of lymphocytes forming nonimmune rosettes with sheep red blood cells, 207
— resorption-stimulating factor produced by mouse fibrosarcoma cells, evidence is prostaglandin E_2, new model for hypercalcemia of cancer, 1329
Bordetella pertussis, distribution of labeled lymph node cells during lymphocytosis induced by, 1581
Bovine serum albumin, methylated, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against, 1616
Brain, catecholamine metabolism, effects of neonatal infection, perinatal malnutrition, and crowding on, lasting biological effects of early environmental influences, 1031
— protein, cellular immunity to, and disease production, dissociation in allergic encephalitis, 156
Brain-associated stem cell antigen shared by brain and hemopoietic stem cells, 369
Breeding, closely bred colony of dogs, histocompatibility studies, tolerance to bone marrow, kidney, and skin allografts in DL-A-identical radiation chimeras, 1080
Bursa cells, chick embryo, synthesis of primordial immunoglobulin M by, 8
Bursal function, ontogeny, immunocompetent cell for humoral immunity, 816

CANCER, hypercalcemia, new model for, evidence that bone resorption-stimulating factor produced by mouse fibrosarcoma cells is prostaglandin E_2, 1329
Carcinoembryonic antigen activity, structural studies of major glycoprotein in preparations with, 200
Carrier (mouse γ-globulin), isologous, immunity and tolerance to hapten (NIP) coupled to, 546
—, nonimmunogenic, immune response to hapten coupled to, influence of lipopolysaccharide, 392
— priming, regulation of hapten-specific antibody class by, 1661
Catecholamine metabolism, brain, effects of neonatal infection, perinatal malnutrition, and crowding on, lasting biological effects of early environmental influences, 1031
Cell antigen, stem, brain-associated, shared by brain and hemopoietic stem cells, 369
— antigens, chemically altered, selective priming of thymus-derived cells by, 656
—, bone marrow-derived, activation in vivo by non-antigen-specific interaction with thymus-derived cells, frequency of immune responses increased by immunization with two antigens, 381
— cooperation mechanism and immunological adjuvants, 185
—, immunocompetent, for humoral immunity, ontogeny of bursal function, 816

Cell immunoglobulin, thymus-derived, and antigen complexes, specific collaboration via in immune response in vitro, 737
— immunoglobulins, thymus-derived (T), presence of receptor site for IgG and absence of large amounts of buried Ig determinants on T cells, 1323
— interactions in immune response in vitro, comparison of effects of antigen-specific and allogeneic thymus-derived cell factors, 722
— interactions in immune response in vitro, specific collaboration across cell-impermeable membrane, 49
— interactions in immune response in vitro, specific collaboration via complexes of antigen and thymus-derived cell immunoglobulin, 737
— lines, lymphoid, cultured, study, Hurler syndrome, 644
— proliferation, thymus-independent (B), in spleen cell cultures of radiation chimeras stimulated by phytohemagglutinin or allogeneic cells, 962
—, responding, in potentiation of thymus-processed lymphocyte response to mitogens, 128
— responses, non-thymus-processed (B) and thymus-processed (T), against methylated bovine serum albumin, analysis of, 1616
—, stem, differentiation, and antibody formation, regulatory effects of sex hormones on, 1098
—, stimulating, in syngeneic and allogeneic mixed lymphocyte reaction, nature of, 1648
— surface immunoglobulin, distribution among thymocytes, bone marrow cells, and derived populations, 81
— surface immunoglobulin, release from murine splenic lymphocytes, 676
— suspensions, spleen, inhibitory and stimulatory effects of concanavalin A on response to antigen, characterization of inhibitory cell activity, 1445
Cell, thymus-derived, activation by heterologous antigens as replacement of specific immune T cells in transfer of secondary response to erythrocytes, 715
—, tumor, surface antigen, virally determined, in vitro cytotoxicity by non-thymus-processed lymphocyte population with specificity for, 1072
— tumors, plasma, syngeneic murine, immune and natural antibodies to, 94
Cell-bound and subcellular alloantigens, requirement for macrophages in cytotoxic reactions against, 331
Cell-host response in local graft-*versus*-host reaction induced in kidneys of F_1 rats by parental thoracic duct lymphocytes, 21
Cell-mediated immune response in vitro, requirement for macrophages in cytotoxic reactions against cell-bound and subcellular alloantigens, 331
— immunity, depressed, in patients with primary intracranial tumors, characterization of humoral immunosuppressive factor, 1631
— tumor destruction in vitro by peritoneal exudate lymphoid cells, pathway for, 1594
Cells, allogeneic, or phytohemagglutinin, thymus-independent (B) cell proliferation in spleen cell cultures of radiation chimeras stimulated by, 962
—, antibody-forming, evidence for intracellular mechanism of inactivation of hapten-specific precursors of, immunological tolerance in bone marrow-derived lymphocytes, 1404
—, blood, red. *See* Erythrocytes.
—, bone marrow-derived and thymus-derived, role in immunological memory, in vitro studies, 663
—, bone marrow, thymocytes, and derived populations, cell surface im-

munoglobulin distribution among, 81
Cells, brain and hemopoietic stem, brain-associated stem cell antigen shared by, 369
—, bursa, chick embryo, synthesis of primordial immunoglobulin M by, 8
—, clonal precursor, primary and secondary, mechanism of antigenic stimulation, 241
— expressing histamine receptors, regulation of antibody response by, 1302
—, fibrosarcoma, mouse, evidence that bone resorption-stimulating factor produced by is prostaglandin E_2, new model for hypercalcemia of cancer, 1329
— from aorta, lysosomes of arterial wall, isolation and subcellular fractionation, 1117
—, hapten-binding, deletion by highly radioactive ^{125}I conjugate, 305
—, immune, specific, thymus-derived, T cell activation by heterologous antigens as replacement of in transfer of secondary response to sheep erythrocytes, 715
—, lymph node, labeled, distribution during lymphocytosis induced by *Bordetella pertussis*, 1581
—, lymphoid, peritoneal exudate, pathway for cell-mediated tumor destruction in vitro by, 1594
—, lymphoid, thymus-independent and thymus-derived, immunofluorescent and functional studies, 984
—, lymphoid, thymus-independent and thymus-derived, ultrastructural studies, 1008
—, lymphoma, grafted, antibody-mediated suppression of, participation of macrophages, 375
—, mammalian, and *Toxoplasma gondii*, interaction between, absence of lysosomal fusion with phagocytic vacuoles containing living parasites, 1173
—, mammalian, and *Toxoplasma gondii*, interaction between, mechanism of entry and intracellular fate of parasite, 1157
Cells, memory, B, and virgin thymus-independent (B) lymphocytes, fundamental changes in physiological characteristics, 851
—, non-thymus-dependent, hapten inhibition studies of antigen binding to in immunity and tolerance, 532
—, spleen, donor and host, role in cytotoxicity in graft-*versus*-host reaction, 39
—, spleen, nonadherent, in vitro, effects of mercaptoethanol and peritoneal macrophages on antibody-forming capacity of, 604
—, target, specific adherence of in vitro differentiated lymphocytes to, 1344
—, thymic and bone marrow, of neonatally splenectomized CBA mice, splenic replenishment of synergistic ability to, 761
—, thymus, immunologic properties, identification of T cell functions within minor, low-density subpopulation, 1461
—, thymus-dependent (T), and macrophages, regulatory effect on response of thymus-independent (B) lymphocytes to endotoxin, 956
—, thymus-derived, bone marrow-derived cell activation in vivo by non-antigen-specific interaction with, frequency of immune responses increased by immunization with two antigens, 381
—, thymus-derived, selective priming by chemically altered cell antigens, 656
—, thymus-influenced, response to histocompatibility-2 antigens, dose-response kinetics, 112
— transformed by murine sarcoma virus, lack of distinctive surface antigen on, 344
Cell-to-cell interaction in immune response, regulation of hapten-specific

antibody class by carrier priming, 1661
Cellular contribution, host, immune lymphocyte transfer reactions with special reference to, 1545
— events during induction of thyroiditis in rabbit, 1605
— immune response to Gross virus-induced lymphoma, in vitro activation of, 969
— immunity to brain protein and disease production, dissociation in allergic encephalitis, 156
— interactions required for antibody production enhancement by graft-*versus*-host reaction, establishment of, 455
— receptors, binding of radioactively labeled anti-alloantiserum, 872
— source of potentiating mediator in potentiation of thymus-processed lymphocyte response to mitogens, 143
Chain, side, of tyrosine, structural requirements in for immunogenicity of L-tyrosine-azobenzenearsonate, 387
Chemically altered cell antigens, selective priming of thymus-derived cells by, 656
— modified and native albumins, immune responses against, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against methylated bovine serum albumin, 1616
Chimeras, radiation, DL-A-identical, tolerance to bone marrow, kidney, and skin allografts in, histocompatibility studies in closely bred colony of dogs, 1080
—, radiation, origin of alveolar macrophages in, 630
—, radiation, stimulated by phytohemagglutinin or allogeneic cells, thymus-independent (B) cell proliferation in spleen cell cultures of, 962
Cholinergic and α-adrenergic stimulation, enhancement of immunological release of histamine and slow-reacting substance of anaphylaxis from human lung by, 556
Ciliary inhibition, oyster, by cystic fibrosis culture medium, 1313
Circulation, micro-, blood, in lymph node during primary immune response, 697
Clonal precursor cells, primary and secondary, mechanism of antigenic stimulation, 241
Cobra venom-induced hemolysis, activity levels in sera of patients with neoplastic and other diseases, 950
Colchicine and vinblastine, effect on topographical separation of membrane functions, 1
Complement-dependent platelet injury by staphylococcal protein A, 68
Concanavalin A, inhibitory and stimulatory effects on response of spleen cell suspensions to antigen, characterization of inhibitory cell activity, 1445
Conjugate, ^{125}I, highly radioactive, deletion of hapten-binding cells by, 305
Conjugates, hapten protein, induction of immunity and tolerance in vitro by, 532
Crosses with *Fv-1*^{b} strains of mice, genetic transmission of murine leukemia virus by AKR mice, 1286
— with *Fv-1*^{n} strains of mice, genetic transmission of murine leukemia virus by AKR mice, 1272
Crowding, neonatal infection, and perinatal malnutrition, effects on catecholamine metabolism of brain, lasting biological effects of early environmental influences, 1031
Culture, leukocyte, mixed, and graft-*versus*-host reaction, reactivity of lymphocytes from germfree rats in, 417
—, leukocyte, mixed, responsiveness of germfree mice in, reexamining nature of responding unit, 398

Culture medium, cystic fibrosis, oyster ciliary inhibition by, 1313
Cultured lymphoid cell lines, study, Hurler syndrome, 644
— skin fibroblasts, deficiency of α-L-fucosidase in, 197
Cultures, spleen cell, of radiation chimeras stimulated by phytohemagglutinin or allogeneic cells, thymus-independent (B) cell proliferation in, 962
Cyclophosphamide, effects on mechanism of recovery from systemic vaccinia virus infection, 277
Cystic fibrosis culture medium, oyster ciliary inhibition by, 1313
Cytotoxic reactions against cell-bound and subcellular alloantigens, requirement for macrophages in, 331
Cytotoxicity assays, idiotypic determinants of immunoglobulin M detected on surface of human lymphocytes by, 650
— in graft-*versus*-host reaction, role of donor and host spleen cells, 39
—, in vitro, by non-thymus-processed lymphocyte population with specificity for virally determined tumor cell surface antigen, 1072

DIETARY potassium and sodium/potassium molar ratios, influence on salt hypertension development, 318
Disease, liver, hereditary, acute intermittent porphyria, defect in reductive transformation of natural steroid hormones in, 1043
— production and cellular immunity to brain protein, dissociation in allergic encephalitis, 156
Diseases, neoplastic and other, cobra venom-induced hemolysis activity levels in sera of patients with, 950
Dose-response kinetics, response of thymus-influenced cells to histocompatibility-2 antigens, 112
Duct lymphocytes, thoracic, parental, host-cell response in local graft-*versus*-host reaction induced in kidneys of F_1 rats by, 21
Duct, thoracic, lymphocytes, lysosomes in, 1394

EGG white lysozyme, cross-tolerance between serologically noncross-reacting forms of, 1308
Electron microscope study of fibrous plaque formation, arteriosclerosis, in primates, 769
Electron-opaque tracers, evidence for blood-thymus barrier using, 466
Embryo, chick, synthesis of primordial immunoglobulin M by bursa cells of, 8
Encephalitis, allergic, dissociation of cellular immunity to brain protein and disease production in, 156
—, viral, specificity of inflammatory response in, adoptive immunization of immunosuppressed mice infected with Sindbis virus, 216
Endothelium, vascular, in lymphoid tissue, migration of lymphocytes across, reexamination, 568
Endotoxin, regulatory effect of macrophages and thymus-dependent (T) cells on response of thymus-independent (B) lymphocytes to, 956
Environmental influences, early, lasting biological effects, effects of neonatal infection, perinatal malnutrition, and crowding on catecholamine metabolism of brain, 1031
Erythrocyte antibody response, anti-sheep, and immunoglobulin synthesis in backcross and F_2 progeny of mice genetically selected for high or low antibody synthesis, genetic factors controlling, 790
— membranes, human, preparation and further characterization of MN glycoprotein of, 1140
Erythrocytes, sheep, induction of tolerance to, 1666
—, sheep, large population of human thymus-derived and bone marrow-

derived lymphocytes forming non-immune rosettes with, 207
Erythrocytes, sheep, thymus-derived cell, cell activation by heterologous antigens as replacement of specific immune T cells in transfer of secondary response to, 715
—, sheep, variations in responses of C57BL/10J and A/J mice to, serological characterization and genetic analysis, 1063
Exudate, peritoneal, lymphoid cells, pathway for cell-mediated tumor destruction in vitro by, 1594

FEVER, penicillin-induced, mechanism, 227
Fibrinolytic pathway of human plasma, isolation and characterization of plasminogen proactivator, 1378
Fibroblasts, skin, cultured, deficiency of α-L-fucosidase in, 197
Fibrosarcoma cells, mouse, evidence that bone resorption-stimulating factor produced by is prostaglandin E_2, new model for hypercalcemia of cancer, 1329
Fibrosis, cystic, culture medium, oyster ciliary inhibition by, 1313
Fibrous plaque formation, arteriosclerosis, in primates, electron microscope study, 769
Fluorescence, immuno-, and ultrastructural radioautography, analysis of ligand-induced movement of lymphocyte membrane macromolecules by, 885
Fluorescent, immuno-, and functional studies of thymus-independent and thymus-derived lymphoid cells, 984
Freeze-etch, freeze-fracture studies on gonococci, 1258
Freeze-fracture, freeze-etch studies on gonococci, 1258
Friend spleen focus-forming virus, neutralization of pseudotypes by type-specific murine antisera, 832
α-L-Fucosidase, deficiency in cultured skin fibroblasts, fucosidosis, 197
Fucosidosis, deficiency of α-L-fucosidase in cultured skin fibroblasts, 197

β-D-GALACTOSIDASE-ACTIVATING antibody, anti-, focal fluctuation and changes in affinity of, 353
Gene function, immune response, histocompatibility-linked, specific inhibition of antigen-induced lymphocyte proliferation by alloantisera, 1207
—, X-linked, evidence for decisive role in determining responsiveness to Type III pneumococcal polysaccharide, 931
Genes, immune response (Ir), H-2-linked, independent loci for *Ir-IgG* and *Ir-IgA* genes, 1231
Genetic analysis and serological characterization of variations in responses of C57BL/10J and A/J mice to sheep red blood cells, 1063
— control of antibody response to Type III pneumococcal polysaccharide, evidence that X-linked gene plays decisive role in determining responsiveness, 931
— control of immune response, effect of graft-*versus*-host reaction on antibody response to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys, 1195
— control of immune responsiveness to IgG (γ2a) allotypes in mouse, association of H-2 types with, 1222
— control of in vivo-in vitro discrepancies in 19S antibody synthesis, immune responses in vitro, 1241
— factors controlling anti-sheep erythrocyte antibody response and immunoglobulin synthesis in backcross and F_2 progeny of mice genetically selected for high or low antibody synthesis, 790
— factors influencing C-type RNA virus induction, 175

Genetic nonresponder mice, antigen binding and capping by lymphocytes of, 403
— transmission of murine leukemia virus by AKR mice, crosses with *Fv-1*^{b} strains of mice, 1286
— transmission of murine leukemia virus by AKR mice, crosses with *Fv-1*^{n} strains of mice, 1272
Germfree mice, responsiveness in mixed leukocyte culture, reexamining nature of responding unit, 398
— rats, lymphocytes from, reactivity in mixed leukocyte culture and in graft-*versus*-host reaction, 417
γ-Globulin, aggregated, interaction with B lymphocytes, 191
—, mouse, isologous carrier, immunity and tolerance to hapten (NIP) coupled to, 546
Glucosamine incorporation by peritoneal macrophages in delayed hypersensitivity, antigen-induced stimulation of, 1518
Glycoprotein, major, in preparations with carcinoembryonic antigen activity, structural studies of, 200
—, MN, of human erythrocyte membranes, preparation and further characterization, 1140
Gonococci, freeze-fracture, freeze-etch studies on, 1258
Gonococcus infection, freeze-fracture, freeze-etch studies on gonococci, 1258
Graft-*versus*-host reaction, conditions for enhancement of hapten-specific secondary antibody responses by, 439
— reaction, cytotoxicity in, role of donor and host spleen cells, 39
— reaction, effect on antibody response to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys, genetic control of immune response, 1195
— reaction, establishment of cellular interactions required for antibody production enhancement by, 455
Graft-*versus*-host reaction, local, induced in kidneys of F_1 rats by parental thoracic duct lymphocytes, host-cell response in, 21
— reaction and mixed leukocyte culture, reactivity of lymphocytes from germfree rats in, 417
— reaction, rapid viral induction of murine lymphomas in, 1533
Grafted lymphoma cells, antibody-mediated suppression of, participation of macrophages, 375
Gross virus-induced lymphoma, in vitro activation of cellular immune response to, 969

HAPTEN (NIP) coupled to isologous carrier (mouse γ-globulin), immunity and tolerance to, 546
— coupled to nonimmunogenic carrier, immune response to, influence of lipopolysaccharide, 392
— inhibition studies of antigen binding to non-thymus-dependent cells in immunity and tolerance, 532
Hapten-binding cells, deletion by highly radioactive ^{125}I conjugate, 305
Hapten-specific antibody class, regulation by carrier priming, cell-to-cell interaction in immune response, 1661
— precursors of antibody-forming cells, evidence for intracellular mechanism of inactivation of, immunological tolerance in bone marrow-derived lymphocytes, 1404
— secondary antibody responses, conditions for enhancement by graft-*versus*-host reaction, 439
— tolerance, preferential depression of high affinity antibody response, 426
Hemolysis, cobra venom-induced, activity levels in sera of patients with neoplastic and other diseases, 950
Hemopoietic stem cells and brain,

brain-associated stem cell antigen shared by, 369
Hereditary liver disease, acute intermittent porphyria, defect in reductive transformation of natural steroid hormones in, 1043
Heterologous antigens, thymus-derived, cell activation by as replacement of specific immune T cells in transfer of secondary response to erythrocytes, 715
— enhancement, specific, of immune responses, specific generation of thymus-derived enhancing factor, 1501
Histamine receptors, regulation of antibody response by cells expressing, 1302
— release from platelets, leukocyte-dependent, role of IgE, basophils, and platelet-activating factor, 1356
— and slow-reacting substance of anaphylaxis from human lung, immunological release, enhancement by cholinergic and α-adrenergic stimulation, 556
Histocompatibility-2 antigens, response of thymus-influenced cells to, dose-response kinetics, 112
Histocompatibility complex, major, serologically defined and lymphocyte-defined components, 1430
— studied in closely bred colony of dogs, tolerance to bone marrow, kidney, and skin allografts in DL-A-identical radiation chimeras, 1080
Histocompatibility-linked immune response gene function, specific inhibition of antigen-induced lymphocyte proliferation by alloantisera, 1207
Hormones, sex, regulatory effects on antibody formation and stem cell differentiation, 1098
—, steroid, natural, in hereditary liver disease, acute intermittent porphyria, defect in reductive transformation of, 1043
Host-*versus*-graft reaction, conditions for enhancement of hapten-specific secondary antibody responses by, 439
— reaction, cytotoxicity in, role of donor and host spleen cells, 39
— reaction, effect on antibody response to poly-L(Tyr,Glu)-poly-D, L-Ala--poly-L-Lys, genetic control of immune response, 1195
— reaction, establishment of cellular interactions required for antibody production enhancement by, 455
— reaction, local, induced in kidneys of F_1 rats by parental thoracic duct lymphocytes, host-cell response in, 21
— reaction and mixed leukocyte culture, reactivity of lymphocytes from germfree rats in, 417
— reaction, rapid viral induction of murine lymphomas in, 1533
Humoral immunity, immunocompetent cell for, ontogeny of bursal function, 816
— immunosuppressive factor, characterization in depressed cell-mediated immunity in patients with primary intracranial tumors, 1631
Hurler syndrome, study of cultured lymphoid cell lines, 644
Hypercalcemia of cancer, new model for, evidence that bone resorption-stimulating factor produced by mouse fibrosarcoma cells is prostaglandin E_2, 1329
Hyperimmunization with streptococci, occurrence of 19S and 7S anti-IgGs during, 799
Hypersensitivity, delayed, antigen-induced stimulation of glucosamine incorporation by peritoneal macrophages in, 1518
Hypertension, salt, influence of dietary potassium and sodium/potassium molar ratios on development of, 318

IDIOTYPIC determinants of immunoglobulin M detected on

surface of human lymphocytes by cytotoxicity assays, 650
Immune cells, specific, thymus-derived, T cell activation by heterologous antigens as replacement of in transfer of secondary response to sheep erythrocytes, 715
— complexes, soluble, in vitro adherence to macrophages, 514
— lymphocyte transfer reactions, with special reference to cellular contribution of host, 1545
— and natural antibodies to syngeneic murine plasma cell tumors, 94
—, non-, rosettes, large population of lymphocytes forming with sheep red blood cells, surface markers on human thymus-derived and bone marrow-derived lymphocytes, 207
— response and antigen recognition, self-help with symmetrical bifunctional antigen molecules, 1478
— response and antigen recognition, structural requirements in tyrosine side chain for immunogenicity of L-tyrosine-azobenzenearsonate, 387
— response, cell-mediated, in vitro, requirement for macrophages in cytotoxic reactions against cell-bound and subcellular alloantigens, 331
— response, cell-to-cell interaction in, regulation of hapten-specific antibody class by carrier priming, 1661
— response, cellular, to Gross virus-induced lymphoma, in vitro activation of, 969
— response gene function, histocompatibility-linked, specific inhibition of antigen-induced lymphocyte proliferation by alloantisera, 1207
— response (Ir) genes, H-2-linked, independent loci for *Ir-IgG* and *Ir-IgA* genes, 1231
— response, genetic control, effect of graft-*versus*-host reaction on antibody response to poly-L(Tyr, Glu)-poly-D, L-Ala--poly-L-Lys, 1195
Immune response in vitro, cell interactions in, comparison of effects of antigen-specific and allogeneic thymus-derived cell factors, 722
— response in vitro, cell interactions in, specific collaboration across cell-impermeable membrane, 49
— response in vitro, cell interactions in, specific collaboration via complexes of antigen and thymus-derived cell immunoglobulin, 737
— response in vitro, maturation, focal fluctuation and changes in affinity of anti-β-D-galactosidase-activating antibody, 353
— response, primary, blood microcirculation in lymph node during, 697
— response to hapten coupled to non-immunogenic carrier, influence of lipopolysaccharide, 392
— responses against native and chemically modified albumins, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against methylated bovine serum albumin, 1616
— responses, frequency increased by immunization with two antigens, bone marrow-derived cell activation in vivo by non-antigen-specific interaction with thymus-derived cells, 381
— responses in vitro, genetic control of in vivo-in vitro discrepancies in 19S antibody synthesis, 1241
— responses, specific heterologous enhancement, specific generation of thymus-derived enhancing factor, 1501
— responsiveness to IgG (γ2a) allotypes in mouse, association of H-2 types with genetic control of, 1222
Immunity, cell-mediated, depressed, in patients with primary intracranial tumors, characterization of humoral immunosuppressive factor, 1631

Immunity, cellular, to brain protein, and disease production, dissociation in allergic encephalitis, 156
—, humoral, immunocompetent cell for, ontogeny of bursal function, 816
— and tolerance, induction in vitro by hapten protein conjugates, 532
— and tolerance to hapten (NIP) coupled to isologous carrier (mouse γ-globulin), 546
Immunization, adoptive, of immunosuppressed mice infected with Sindbis virus, specificity of inflammatory response in viral encephalitis, 216
— with two antigens, frequency of immune responses increased by, bone marrow-derived cell activation in vivo by non-antigen-specific interaction with thymus-derived cells, 381
Immunocompetent cell for humoral immunity, ontogeny of bursal function, 816
— cells, surface antigens, in vitro studies of role of bone marrow-derived and thymus-derived cells in immunological memory, 663
Immunofluorescence and ultrastructural radioautography, analysis of ligand-induced movement of lymphocyte membrane macromolecules by, 885
Immunofluorescent and functional studies of thymus-independent and thymus-derived lymphoid cells, 984
Immunogenic, non-, carrier, immune response to hapten coupled to, influence of lipopolysaccharide, 392
Immunogenicity of L-tyrosine-azobenzenearsonate, structural requirements in tyrosine side chain for, 387
Immunoglobulin, cell surface, distribution among thymocytes, bone marrow cells, and derived populations, 81
, cell surface, release from murine splenic lymphocytes, 676
Immunoglobulin, cell, thymus-derived, and antigen complexes, specific collaboration via in immune response in vitro, 737
— M detected on surface of human lymphocytes by cytotoxicity assays, idiotypic determinants, 650
— M, primordial, synthesis by chick embryo bursa cells, 8
— synthesis and anti-sheep erythrocyte antibody response in backcross and F_2 progeny of mice genetically selected for high or low antibody synthesis, genetic factors controlling, 790
Immunoglobulins, cell, thymus-derived (T), presence of receptor site for IgG and absence of large amounts of buried Ig determinants on T cells, 1323
Immunologic properties of thymus cells, identification of T cell functions within minor, low-density subpopulation, 1461
Immunological adjuvants and cell cooperation mechanism, 185
— memory, in vitro studies of role of bone marrow-derived and thymus-derived cells in, 663
— release of histamine and slow-reacting substance of anaphylaxis from human lung, enhancement by cholinergic and α-adrenergic stimulation, 556
— tolerance in bone marrow-derived lymphocytes, evidence for intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404
Immunologically specific macrophage-arming factor produced by thymus-derived lymphocytes, 1318
Immunosuppressed mice infected with Sindbis virus, adoptive immunization of, specificity of inflammatory response in viral encephalitis, 216
Immunosuppressive factor, humoral, characterization in depressed cell-mediated immunity in patients with primary intracranial tumors, 1631

Inbred mice, allogeneic effect in, conditions for enhancement of hapten-specific secondary antibody responses by graft-*versus*-host reaction, 439
— mice, allogeneic effect in, establishment of cellular interactions required for antibody production enhancement by graft-*versus*-host reaction, 455
Infection, gonococcus, freeze-fracture, freeze-etch studies on gonococci, 1258
—, neonatal, perinatal malnutrition, and crowding, effects on catecholamine metabolism of brain, lasting biological effects of early environmental influences, 1031
—, vaccinia virus, systemic, mechanism of recovery from, effects of cyclophosphamide, 277
Inflammatory response in viral encephalitis, specificity, adoptive immunization of immunosuppressed mice infected with Sindbis virus, 216
Inhibition, ciliary, oyster, by cystic fibrosis culture medium, 1313
—, specific, of antigen-induced lymphocyte proliferation by alloantisera, histocompatibility-linked immune response gene function, 1207
— studies, hapten, of antigen, binding to non-thymus-dependent cells in immunity and tolerance, 532
Inhibitory factor, migration (MIF), evidence for receptor present on peritoneal macrophage but not on alveolar macrophage, 589
Injury, platelet, complement-dependent, by staphylococcal protein A, 68
Inoculation with material from multiple sclerosis patients, decreased percentage of polymorphonuclear neutrophils in peripheral blood after, 618
Intracellular fate of parasite and mechanism of entry, interaction between *Toxoplasma gondii* and mammalian cells, 1157
— mechanism of inactivation of hapten-specific precursors of antibody-forming cells, evidence for, immunological tolerance in bone marrow-derived lymphocytes, 1404
Intracranial tumors, primary, depressed cell-mediated immunity in patients with, characterization of humoral immunosuppressive factor, 1631
Iodine-125 conjugate, highly radioactive, deletion of hapten-binding cells by, 305
Isologous carrier, mouse γ-globulin, immunity and tolerance to hapten (NIP) coupled to, 546

KIDNEY, bone marrow, and skin allografts in DL-A-identical radiation chimeras, tolerance to, histocompatibility studies in closely bred colony of dogs, 1080
—, protein transport in utilizing tyrosinase as ultrastructural tracer, 291
Kidneys of F_1 rats, host-cell response in local graft-*versus*-host reaction induced by parental thoracic duct lymphocytes in, 21
Kinetics, dose-response, response of thymus-influenced cells to histocompatibility-2 antigens, 112

LEUKEMIA virus, murine, genetic transmission by AKR mice, crosses with *Fv-1*^{b} strains of mice, 1286
— virus, murine, genetic transmission by AKR mice, crosses with *Fv-1*^{n} strains of mice, 1272
— viruses, murine, classification, neutralization of pseudotypes of Friend spleen focus-forming virus by type-specific murine antisera, 832
Leukocyte culture, mixed, and graft-*versus*-host reaction, reactivity of lymphocytes from germfree rats in, 417
— culture, mixed, responsiveness of germfree mice in, reexamining nature of responding unit, 398
Leukocyte-dependent histamine release from platelets, role of IgE,

basophils, and platelet-activating factor, 1356
Leukocytes, human, neutrophil-immobilizing factor derived from, generation and partial characterization, 1564
Lipopolysaccharide, influence on immune response to hapten coupled to nonimmunogenic carrier, 392
Liver disease, hereditary, acute intermittent porphyria, defect in reductive transformation of natural steroid hormones in, 1043
Lung, human, immunological release of histamine and slow-reacting substance of anaphylaxis from, enhancement by cholinergic and α-adrenergic stimulation, 556
Lymph node, blood microcirculation in during primary immune response, 697
— node cells, labeled, distribution during lymphocytosis induced by *Bordetella pertussis*, 1581
Lymphocyte, immune, transfer reactions, with special reference to cellular contribution of host, 1545
— membrane macromolecules, ligand-induced movement, analysis by immunofluorescence and ultrastructural radioautography, 885
— membrane macromolecules, ligand-induced movement, mapping of surface moieties, 907
— population, non-thymus-processed, with specificity for virally determined tumor cell surface antigen, in vitro cytotoxicity by, 1072
— populations, thymic, functional maturation in vitro, 1484
— proliferation, antigen-induced, specific inhibition by alloantisera, histocompatibility-linked immune response gene function, 1207
— reaction, syngeneic and allogeneic mixed, nature of stimulating cell in, 1648
— responses in vitro, regulation, 956
—, thymus-processed, potentiation of response to mitogens, cellular source of potentiating mediator, 143
Lymphocyte, thymus-processed, potentiation of response to mitogens, responding cell, 128
Lymphocyte-defined and serologically defined components of major histocompatibility complex, 1430
Lymphocytes, bone marrow-derived, immunological tolerance in, evidence for intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404
— from germfree rats, reactivity in mixed leukocyte culture and in graft-*versus*-host reaction, 417
—, human bone marrow-derived, class, subclass, and allelic exclusion of membrane-bound Ig of, 409
—, human, idiotypic determinants of immunoglobulin M detected on surface by cytotoxicity assays, 650
—, human thymus-derived and bone marrow-derived, large population, forming nonimmune rosettes with red blood cells, surface markers on, 207
—, in vitro differentiated, specific adherence to target cells, 1344
—, B, interaction of aggregated γ-globulin with, 191
—, migration across vascular endothelium in lymphoid tissue, reexamination, 568
— of genetic nonresponder mice, antigen binding and capping by, 403
—, recirculating, specific selection by antigen, 499
—, splenic, murine, release of cell surface immunoglobulin from, 676
—, thoracic duct, lysosomes in, 1394
—, thoracic duct, parental, host-cell response in local graft-*versus*-host reaction induced in kidneys of F_1 rats by, 21
—, thymus-derived, produce immunologically specific macrophage-arming factor, 1318
—, thymus-independent (B), regula-

tory effect of macrophages and thymus-dependent (T) cells on response to endotoxin, 956
Lymphocytes, thymus-independent (B), virgin, and B memory cells, fundamental changes in physiological characteristics, 851
Lymphocytosis induced by *Bordetella pertussis*, distribution of labeled lymph node cells during, 1581
Lymphoid cell lines, cultured, study, Hurler syndrome, 644
— cells, peritoneal exudate, pathway for cell-mediated tumor destruction in vitro by, 1594
— cells, thymus-independent and thymus-derived, immunofluorescent and functional studies, 984
— cells, thymus-independent and thymus-derived, ultrastructural studies, 1008
— system, chick embryo bursa cells, synthesis of primordial immunoglobulin M by, 8
— tissue, migration of lymphocytes across vascular endothelium in, reexamination, 568
Lymphoma cells, grafted, antibody-mediated suppression of, participation of macrophages, 375
—, virus-induced, Gross, in vitro activation of cellular immune response to, 969
Lymphomas, murine, rapid viral induction in graft-*versus*-host reaction, 1533
Lysosomal fusion with phagocytic vacuoles containing living parasites, absence in interaction between *Toxoplasma gondii* and mammalian cells, 1173
Lysosomes in thoracic duct lymphocytes, 1394
— of arterial wall, isolation and subcellular fractionation of cells from aorta, 1117
Lysozyme, egg white, cross-tolerance between serologically non-cross-reacting forms of, 1308

MACROMOLECULES, membrane, lymphocyte, ligand-induced movement, analysis by immunofluorescence and ultrastructural radioautography, 885
Macromolecules, membrane, lymphocyte, ligand-induced movement, mapping of surface moieties, 907
Macrophage, peritoneal, evidence for receptor for migration inhibitory factor (MIF) present on but not on alveolar macrophage, 589
Macrophage-arming factor, immunologically specific, produced by thymus-derived lymphocytes, 1318
Macrophages, alveolar, origin in radiation chimeras, 630
—, in vitro adherence of soluble immune complexes to, 514
—, participation in antibody-mediated suppression of grafted lymphoma cells, 375
—, peritoneal, in delayed hypersensitivity, antigen-induced stimulation of glucosamine incorporation by, 1518
—, peritoneal, and mercaptoethanol, effects on antibody-forming capacity of nonadherent spleen cells in vitro, 604
—, requirement for in cytotoxic reactions against cell-bound and subcellular alloantigens, 331
— and thymus-dependent (T) cells, regulatory effect on response of thymus-independent (B) lymphocytes to endotoxin, 956
Malnutrition, perinatal, neonatal infection, and crowding, effects on catecholamine metabolism of brain, lasting biological effects of early environmental influences, 1031
Mammalian cells and *Toxoplasma gondii*, interaction between, absence of lysosomal fusion with phagocytic vacuoles containing living parasites, 1173
— cells and *Toxoplasma gondii*, interaction between, mechanism of entry and intracellular fate of parasite, 1157
Man, bone marrow-derived lymphocytes, class, subclass, and allelic ex-

clusion of membrane-bound Ig of, 409
Man, erythrocyte membranes, preparation and further characterization of MN glycoprotein of, 1140
—, leukocytes, neutrophil-immobilizing factor derived from, generation and partial characterization, 1564
—, lung, immunological release of histamine and slow-reacting substance of anaphylaxis from, enhancement by cholinergic and α-adrenergic stimulation, 556
—, lymphocytes, idiotypic determinants of immunoglobulin M detected on surface by cytotoxicity assays, 650
—, plasma, fibrinolytic pathway, isolation and characterization of plasminogen proactivator, 1378
—, thymus-derived and bone marrow-derived lymphocytes, large population, forming nonimmune rosettes with sheep red blood cells, surface markers on, 207
Mapping of surface moieties in ligand-induced movement of lymphocyte membrane macromolecules, 907
Marrow, bone, kidney, and skin allografts in DL-A-identical radiation chimeras, tolerance to, histocompatibility studies in closely bred colony of dogs, 1080
—, bone, and thymic cells of neonatally splenectomized CBA mice, splenic replenishment of synergistic ability to, 761
— cells, bone, thymocytes, and derived populations, cell surface immunoglobulin distribution among, 81
Marrow-derived, bone, cell activation in vivo by non-antigen-specific interaction with thymus-derived cells, frequency of immune responses increased by immunization with two antigens, 381
—, bone, lymphocytes, immunological tolerance in, evidence for intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404
Marrow-derived, bone, and thymus-derived cells, role in immunological memory, in vitro studies, 663
—, bone, and thymus-derived lymphocytes, human, surface markers on, large population of lymphocytes forming nonimmune rosettes with sheep red blood cells, 207
Maturation, functional, of thymic lymphocyte populations in vitro, 1484
— of immune response in vitro, focal fluctuation and changes in affinity of anti-β-D-galactosidase-activating antibody, 353
Medium, culture, cystic fibrosis, oyster ciliary inhibition by, 1313
Membrane, cell-impermeable, specific collaboration across, cell interactions in immune response in vitro, 49
— functions, effect of colchicine and vinblastine on topographical separation of, 1
— macromolecules, lymphocyte, ligand-induced movement, analysis by immunofluorescence and ultrastructural radioautography, 885
— macromolecules, lymphocyte, ligand-induced movement, mapping of surface moieties, 907
Membrane-bound Ig of human bone marrow-derived lymphocytes, class, subclass, and allelic exclusion of, 409
Membranes, erythrocyte, human, preparation and further characterization of MN glycoprotein of, 1140
Memory cells, B, and virgin thymus independent (B) lymphocytes, fundamental changes in physiological characteristics, 851
—, immunological, in vitro studies of role of bone marrow-derived and thymus-derived cells in, 663
Mercaptoethanol and peritoneal macrophages, effects on antibody-

forming capacity of nonadherent spleen cells in vitro, 604
Metabolism, catecholamine, brain, effects of neonatal infection, perinatal malnutrition, and crowding on, lasting biological effects of early environmental influences, 1031
Methylated bovine serum albumin, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against, 1616
Microcirculation, blood, in lymph node during primary immune response, 697
Microscope, electron, study of fibrous plaque formation, arteriosclerosis, in primates, 769
Mitogens, potentiation of thymus-processed lymphocyte response to, cellular source of potentiating mediator, 143
-, potentiation of thymus-processed lymphocyte response to, responding cell, 128
Molecules, antigen, symmetrical bifunctional, self-help with, antigen recognition and immune response, 1478
Multiple sclerosis patients, decreased percentage of polymorphonuclear neutrophils in peripheral blood after inoculation with material from, 618

NEONATAL infection, perinatal malnutrition, and crowding, effects on catecholamine metabolism of brain, lasting biological effects of early environmental influences, 1031
Neonatally splenectomized CBA mice, splenic replenishment of synergistic ability to bone marrow and thymic cells of, 761
Neoplastic and other diseases, cobra venom-induced, hemolysis activity levels in sera of patients with, 950
Neovascularization, tumor dormancy in vivo by prevention of, 261
Neutrophil-immobilizing factor derived from human leukocytes, generation and partial characterization, 1564
Neutrophils, polymorphonuclear, decreased percentage in peripheral blood after inoculation with material from multiple sclerosis patients, 618
Node, lymph, blood microcirculation in during primary immune response, 697
—, lymph, cells, labeled, distribution during lymphocytosis induced by *Bordetella pertussis*, 1581
Nucleic acid, ribo-, C-type, virus induction, genetic factors influencing, 175

ONTOGENY of bursal function, immunocompetent cell for humoral immunity, 816
Oyster ciliary inhibition by cystic fibrosis culture medium, 1313

PARASITE, mechanism of entry and intracellular fate, interaction between *Toxoplasma gondii* and mammalian cells, 1157
Parasites, living, absence of lysosomal fusion with phagocytic vacuoles containing, in interaction between *Toxoplasma gondii* and mammalian cells, 1173
Parental thoracic duct lymphocytes, host-cell response in local graft-*versus*-host reaction induced in kidneys of F_1 rats by, 21
Penicillin-induced fever, mechanism, 227
Perinatal malnutrition, neonatal infection, and crowding, effects on catecholamine metabolism of brain, lasting biological effects of early environmental influences, 1031
Peritoneal exudate lymphoid cells, pathway for cell-mediated tumor destruction in vitro by, 1594
— macrophage, evidence for receptor for migration inhibitory factor (MIF) present on but not on alveolar macrophage, 589
— macrophages, in delayed hyper-

sensitivity, antigen-induced stimulation of glucosamine incorporation by, 1518
Peritoneal macrophages and mercaptoethanol, effects on antibody-forming capacity of nonadherent spleen cells in vitro, 604
Phagocytic vacuoles containing living parasites, absence of lysosomal fusion with in interaction between *Toxoplasma gondii* and mammalian cells, 1173
Physiological characteristics of virgin thymus-independent (B) lymphocytes and B memory cells, fundamental changes in, 851
Phytohemagglutinin or allogeneic cells, thymus-independent (B) cell proliferation in spleen cell cultures of radiation chimeras stimulated by, 962
Plaque formation, fibrous, arteriosclerosis, in primates, electron microscope study, 769
Plasma cell tumors, syngeneic murine, immune and natural antibodies to, 94
—, human, fibrinolytic pathway, isolation and characterization of plasminogen proactivator, 1378
Plasminogen proactivator, isolation and characterization, fibrinolytic pathway of human plasma, 1378
Platelet injury, complement-dependent, by staphylococcal protein A, 68
Platelet-activating factor, IgE, and basophils, role in leukocyte-dependent histamine release from platelets, 1356
Platelets, leukocyte-dependent histamine release from, role of IgE, basophils, and platelet-activating factor, 1356
Pneumococcal polysaccharide, Type III, genetic control of antibody response to, evidence that X-linked gene plays decisive role in determining responsiveness, 931
Polymorphonuclear neutrophils, decreased percentage in peripheral blood after inoculation with material from multiple sclerosis patients, 618
Polysaccharide, pneumococcal, Type III, genetic control of antibody response to, evidence that X-linked gene plays decisive role in determining responsiveness, 931
Porphyria, acute intermittent, hereditary liver disease, defect in reductive transformation of natural steroid hormones in, 1043
Potassium, dietary, and sodium/potassium molar ratios, influence on salt hypertension development, 318
Potentiation of thymus-processed lymphocyte response to mitogens, cellular source of potentiating mediator, 143
— of thymus-processed lymphocyte response to mitogens, responding cell, 128
Preimmunization, specific selection of recirculating lymphocytes by antigen in, 499
Primates, electron microscope study of fibrous plaque formation in, arteriosclerosis, 769
Priming, carrier, regulation of hapten-specific antibody class by, 1661
—, selective, of thymus-derived cells by chemically altered cell antigens, 656
Probability analysis [Appendix], 568
Prostaglandin E_2, evidence that bone resorption-stimulating factor production by mouse fibrosarcoma cells is, new model for hypercalcemia of cancer, 1329
Protein, brain, cellular immunity to, and disease production, dissociation in allergic encephalitis, 156
— conjugates, hapten, induction of immunity and tolerance in vitro by, 532
— A, staphylococcal, complement-dependent platelet injury by, 68
— transport in kidney utilizing tyro-

sinase as ultrastructural tracer, 291

RADIATION chimeras, DL-A-identical, tolerance to bone marrow, kidney, and skin allografts in, histocompatibility studies in closely bred colony of dogs, 1080

— chimeras, origin of alveolar macrophages in, 630

— chimeras stimulated by phytohemagglutinin or allogeneic cells, thymus-independent (B) cell proliferation in spleen cell cultures of, 962

Radioactive, highly, ^{125}I conjugate, deletion of hapten-binding cells by, 305

Radioactively labeled anti-alloantiserum, binding of, 872

Radioautography, ultrastructural, and immunofluorescence, analysis of ligand-induced movement of lymphocyte membrane macromolecules by, 885

Receptor for migration inhibitory factor (MIF) present on peritoneal macrophage but not on alveolar macrophage, evidence for, 589

— site for IgG, presence, and absence of large amounts of buried Ig determinants on T cells, thymus-derived (T) cell immunoglobulins, 1323

Receptors, cellular, binding of radioactively labeled anti-alloantiserum, 872

—, histamine, regulation of antibody response by cells expressing, 1302

Resorption-stimulating factor, bone, produced by mouse fibrosarcoma cells, evidence is prostaglandin E_2, new model for hypercalcemia of cancer, 1329

Ribonucleic acid, C-type, virus induction, genetic factors influencing, 175

Rosettes, nonimmune, large population of lymphocytes forming with sheep red blood cells, surface markers on human thymus-derived and bone marrow-derived lymphocytes, 207

SALT hypertension, influence of dietary potassium and sodium/potassium molar ratios on development of, 318

Sarcoma virus, murine, lack of distinctive surface antigen on cells transformed by, 344

Sera of patients with neoplastic and other diseases, cobra venom-induced hemolysis, activity levels in, 950

Serological characterization and genetic analysis of variations in responses of C57BL/10J and A/J mice to sheep red blood cells, 1063

Serologically defined and lymphocyte-defined components of major histocompatibility complex, 1430

— non-cross-reacting forms of egg white lysozyme, cross-tolerance between, 1308

Serum albumin, bovine, methylated, analysis of non-thymus-processed (B) and thymus-processed (T) cell responses against, 1616

Sex hormones, regulatory effects on antibody formation and stem cell differentiation, 1098

Sindbis virus, adoptive immunization of immunosuppressed mice infected with, specificity of inflammatory response in viral encephalitis, 216

Skin allografts, bone marrow, and kidney in DL-A-identical radiation chimeras, tolerance to, histocompatibility studies in closely bred colony of dogs, 1080

— fibroblasts, cultured, deficiency of α-L-fucosidase in, 197

Sodium/potassium and dietary potassium molar ratios, influence on salt hypertension development, 318

Soluble immune complexes, in vitro adherence to macrophages, 514
Specific adherence of in vitro differentiated lymphocytes to target cells, 1344
—, antigen-, and allogeneic thymus-derived cell factors, comparison of effects in immune response in vitro, 722
— collaboration across cell-impermeable membrane, cell interactions in immune response in vitro, 49
— collaboration in immune response in vitro via complexes of antigen and thymus-derived cell immunoglobulin, 737
—, hapten-, antibody class, regulation by carrier priming, 1661
—, hapten-, precursors of antibody-forming cells, evidence for intracellular mechanism of inactivation of, immunological tolerance in bone marrow-derived lymphocytes, 1404
— heterologous enhancement of immune responses, specific generation of thymus-derived enhancing factor, 1501
— immune T cells, thymus-derived, cell activation by heterologous antigens as replacement of in transfer of secondary response to sheep erythrocytes, 715
—, immunologically, macrophage-arming factor produced by thymus-derived lymphocytes, 1318
— inhibition of antigen-induced lymphocyte proliferation by alloantisera, histocompatibility-linked immune response gene function, 1207
—, non-antigen-, interaction with thymus-derived cells, bone marrow-derived cell activation in vivo by, frequency of immune responses increased by immunization with two antigens, 381
— selection of recirculating lymphocytes by antigen, 499
—, type-, murine antisera, neutralization of pseudotypes of Friend spleen focus-forming virus by, 832
Specificity for virally determined tumor cell surface antigen, in vitro cytotoxicity by non-thymus-processed lymphocyte population with, 1072
— of inflammatory response in viral encephalitis, adoptive immunization of immunosuppressed mice infected with Sindbis virus, 216
Spleen cell cultures of radiation chimeras stimulated by phytohemagglutinin or allogeneic cells, thymus-independent (B) cell proliferation in, 962
— cell suspensions, inhibitory and stimulatory effects of concanavalin A on response to antigen, characterization of inhibitory cell activity, 1445
— cells, donor and host, role in cytotoxicity in graft-*versus*-host reaction, 39
— cells, nonadherent, in vitro, effects of mercaptoethanol and peritoneal macrophages on antibody-forming capacity of, 604
— focus-forming virus, Friend, neutralization of pseudotypes by type-specific murine antisera, 832
Splenectomized, neonatally, CBA mice, splenic replenishment of synergistic ability to bone marrow and thymic cells of, 761
Splenic lymphocytes, murine, release of cell surface immunoglobulin from, 676
— replenishment of synergistic ability to bone marrow and thymic cells of neonatally splenectomized CBA mice, 761
Staphylococcal protein A, complement-dependent platelet injury by, 68
Stem cell antigen, brain-associated, shared by brain and hemopoietic stem cells, 369
— cell differentiation and antibody formation, regulatory effects of sex hormones on, 1098

Steroid hormones, natural, in hereditary liver disease, acute intermittent porphyria, defect in reductive transformation of, 1043
Stimulation, antigenic, of primary and secondary clonal precursor cells, mechanism, 241
—, antigen-induced, of glucosamine incorporation by peritoneal macrophages in delayed hypersensitivity, 1518
—, cholinergic and α-adrenergic, enhancement of immunological release of histamine and slow-reacting substance of anaphylaxis from human lung by, 556
Stimulatory and inhibitory effects of concanavalin A on response of spleen cell suspensions to antigen, characterization of inhibitory cell activity, 1445
Streptococci, occurrence of 19S and 7S anti-IgGs during hyperimmunization with, 799
Syndrome, Hurler, study of cultured lymphoid cell lines, 644
Synergistic ability, splenic replenishment to bone marrow and thymic cells of neonatally splenectomized CBA mice, 761
Syngeneic and allogeneic mixed lymphocyte reaction, nature of stimulating cell in, 1648
— murine plasma cell tumors, immune and natural antibodies to, 94

TARGET cells, specific adherence of in vitro differentiated lymphocytes to, 1344
Thoracic duct lymphocytes, lysosomes in, 1394
— duct lymphocytes, parental, host-cell response in local graft-*versus*-host reaction induced in kidneys of F_1 rats by, 21
Thymic and bone marrow cells of neonatally splenectomized CBA mice, splenic replenishment of synergistic ability to, 761
— lymphocyte populations, functional maturation in vitro, 1484
Thymocytes, bone marrow cells, and derived populations, cell surface immunoglobulin distribution among, 81
Thymus cells, immunologic properties, identification of T cell functions within minor, low-density subpopulation, 1461
Thymus-blood barrier, evidence for using electron-opaque tracers, 466
Thymus-dependent (T) cells and macrophages, regulatory effect on response of thymus-independent (B) lymphocytes to endotoxin, 956
—, non-, cells, hapten inhibition studies of antigen binding to in immunity and tolerance, 532
Thymus-derived, allogeneic and antigen-specific cell factors, comparison of effects in immune response in vitro, 722
— and bone-marrow derived cells, role in immunological memory, in vitro studies, 663
— and bone marrow-derived lymphocytes, human, surface markers on, large population of lymphocytes forming nonimmune rosettes with sheep red blood cells, 207
— cell, activation by heterologous antigens as replacement of specific immune T cells in transfer of secondary response to erythrocytes, 715
— cell immunoglobulin and antigen complexes, specific collaboration via in immune response in vitro, 737
— (T) cell immunoglobulins, presence of receptor site for IgG and absence of large amounts of buried Ig determinants on T cells, 1323
— cells, bone marrow-derived cell activation in vivo by non-antigen-specific interaction with, frequency of immune responses increased by immunization with two antigens, 381
— cells, selective priming by chemically altered cell antigens, 656
— enhancing factor, specific genera-

tion, specific heterologous enhancement of immune responses, 1501
Thymus-derived lymphocytes produce immunologically specific macrophage-arming factor, 1318
— and thymus-independent lymphoid cells, immunofluorescent and functional studies, 984
— and thymus-independent lymphoid cells, ultrastructural studies, 1008
Thymus-independent (B) cell proliferation in spleen cell cultures of radiation chimeras stimulated by phytohemagglutinin or allogeneic cells, 962
— (B) lymphocytes, regulatory effect of macrophages and thymus-dependent (T) cells on response to endotoxin, 956
— (B) lymphocytes, virgin, and B memory cells, fundamental changes in physiological characteristics of, 851
— and thymus-derived lymphoid cells, immunofluorescent and functional studies, 984
— and thymus-derived lymphoid cells, ultrastructural studies, 1008
Thymus-influenced cells, response to histocompatibility-2 antigens, dose-response kinetics, 112
Thymus-processed lymphocyte, potentiation of response to mitogens, cellular source of potentiating mediator, 143
— lymphocyte, potentiation of response to mitogens, responding cell, 128
—, non-, lymphocyte population with specificity for virally determined tumor cell surface antigen, in vitro cytotoxicity by, 1072
— (T) and non-thymus-processed (B) cell responses against methylated bovine serum albumin, analysis of, 1616
Thyroiditis in rabbit, cellular events during induction of, 1605
Tissue, lymphoid, migration of lymphocytes across vascular endothelium in, reexamination, 568
Tolerance, cross-, between serologically non-cross-reacting forms of egg white lysozyme, 1308
—, hapten-specific, preferential depression of high affinity antibody response, 426
— and immunity, induction in vitro by hapten protein conjugates, 532
— and immunity to hapten (NIP) coupled to isologous carrier (mouse γ-globulin), 546
—, immunological, in bone marrow-derived lymphocytes, evidence for intracellular mechanism of inactivation of hapten-specific precursors of antibody-forming cells, 1404
— to bone marrow, kidney, and skin allografts in DL-A-identical radiation chimeras, histocompatibility studies in closely bred colony of dogs, 1080
— to sheep red blood cells, induction of, 1666
Topographical separation of membrane functions, effect of colchicine and vinblastine on, 1
Toxoplasma gondii and mammalian cells, interaction between, absence of lysosomal fusion with phagocytic vacuoles containing living parasites, 1173
— *gondii* and mammalian cells, interaction between, mechanism of entry and intracellular fate of parasite, 1157
Tracer, ultrastructural, protein transport in kidney utilizing tyrosinase as, 291
Tracers, electron-opaque, evidence for blood-thymus barrier using, 466
Transformation, reductive, of natural steroid hormones in hereditary liver disease, acute intermittent porphyria, defect in, 1043
Transport, protein, in kidney, utilizing tyrosinase as ultrastructural tracer, 291

Tumor cell surface antigen, virally determined, in vitro cytotoxicity by non-thymus-processed lymphocyte population with specificity for, 1072
— destruction in vitro, cell-mediated, by peritoneal exudate lymphoid cells, pathway for, 1594
— dormancy in vivo by prevention of neovascularization, 261
Tumors, intracranial, primary, depressed cell-mediated immunity in patients with, characterization of humoral immunosuppressive factor, 1631
—, plasma cell, syngeneic murine, immune and natural antibodies to, 94
Tyrosinase as ultrastructural tracer, protein transport in kidney utilizing, 291
Tyrosine, side chain, structural requirements in for immunogenicity of L-tyrosine-azobenzenearsonate, 387
L-Tyrosine-azobenzenearsonate, structural requirements in side chain of tyrosine for immunogenicity of, 387

ULTRASTRUCTURAL radioautography and immunofluorescence, analysis of ligand-induced movement of lymphocyte membrane macromolecules by, 885
— studies of thymus-independent and thymus-derived lymphoid cells, 1008
— tracer, protein transport in kidney utilizing tyrosinase as, 291

VACCINIA virus infection, systemic, mechanism of recovery from, effects of cyclophosphamide, 277
Vacuoles, phagocytic, containing living parasites, absence of lysosomal fusion with in interaction between *Toxoplasma gondii* and mammalian cells, 1173
Vascular endothelium in lymphoid tissue, migration of lymphocytes across, reexamination, 568
Venom-induced hemolysis, cobra, activity levels in sera of patients with neoplastic and other diseases, 950
Vinblastine and colchicine, effect on topographical separation of membrane functions, 1
Viral encephalitis, specificity of inflammatory response in, adoptive immunization of immunosuppressed mice infected with Sindbis virus, 216
— induction, rapid, of murine lymphomas in graft-*versus*-host reaction, 1533
Virally determined tumor cell surface antigen, in vitro cytotoxicity by non-thymus-processed lymphocyte population with specificity for, 1072
Virus, Friend spleen focus-forming, neutralization of pseudotypes by type-specific murine antisera, 832
— induction, C-type RNA, genetic factors influencing, 175
— infection, vaccinia, systemic, mechanism of recovery from, effects of cyclophosphamide, 277
—, leukemia, murine, genetic transmission by AKR mice, crosses with *Fv-1*^{b} strains of mice, 1286
—, leukemia, murine, genetic transmission by AKR mice, crosses with *Fv-1*^{n} strains of mice, 1272
—, sarcoma, murine, lack of distinctive surface antigen on cells transformed by, 344
—, Sindbis, adoptive immunization of immunosuppressed mice infected with, specificity of inflammatory response in viral encephalitis, 216
Virus-induced lymphoma, Gross, in vitro activation of cellular immune response to, 969
Viruses, leukemia, murine, classification, neutralization of pseudotypes of Friend spleen focus-forming virus by type-specific murine antisera, 832